

Modular and equitable stable matching rules^{*}

Ahmet Alkan[†]

Kemal Yıldız[‡]

October 13, 2022

Abstract

We propose and analyze *modular stable matching rules* as a candidate for a foundational framework to address issues of social welfare and equity in the stable matching model. We present two characterizations for modular stable matching rules that reveal the ordinal content of optimizing a *modular* function under the *stability* constraint, and present several examples. Then, we propose a new class of equity notions and characterize the class of modular stable matching rules that comply with these notions. Our analysis indicates that modular matching rules are both structured and rich enough to implement a wide range of objectives.

Keywords: Equity, attainability, lattice, rotations, modular optimization.

^{*}We thank Battal Doğan, Serhat Doğan, Faruk Gul, Tarık Kara, Deniz Savas and participants of several seminars and conferences for valuable comments and suggestions. Kemal Yıldız's research is supported by the BAGEP Award of the Science Academy in Turkey. First posted version: February 8, 2022.

[†]Department of Economics, Sabancı University; alkan@sabanciuniv.edu.tr

[‡]Department of Economics, Bilkent University and Princeton University; kemal.yildiz@bilkent.edu.tr.

Contents

1	Introduction	3
2	Modular Stable Matching rules	8
2.1	Stable matching rules	8
2.2	Modularity	11
3	Axioms and characterizations	15
3.1	Convexity	15
3.2	Independence of irrelevant rankings	19
4	A new (class of) equity notions	22
5	Related literature and final comments	24
6	Appendix A: Stepping stones	31
6.1	Rotations and some preliminary observations	31
6.2	An order isomorphism result	36
7	Appendix B	41
7.1	Proof of Theorem 1	41
7.2	Proof of Proposition 2	47
7.3	Proof of Theorem 2	49

8	Appendix C	53
8.1	On the structure of (stable) mixtures	53
8.2	An example	55

1 Introduction

An important feature of two sided matching markets is that there typically exist many stable matchings. These matchings form a lattice according to the group preferences of agents on either side, with the so-called *polarity* property that says if all agents on one side prefer one matching to another than the opposite is true for agents on the other side. Since the pioneering work of [Gale & Shapley \(1962\)](#), attention has mostly rested on the extremal matchings, optimal for one side pessimal for the other, mainly for the remarkable properties they own. The *Deferred Acceptance Procedure* finds the optimal matching very efficiently. The inequity at the extremal matchings, on the other hand, has surely generated interest for compromise solutions but, informational or computational concerns aside, the “middle” has turned out and remained a challenging matter. We believe that existing studies in several disciplines have proven that the breadth of possibilities calls for a foundational framework to address issues of social welfare and equity in the stable matching model. We propose and analyze the *modular stable matching rules* as a candidate to form such a framework mainly due to their analytical tractability along with clarity and richness.

Non-extremal stable matchings have been on the agenda for a long time. There is now a list of solutions that have been put forward, some on geometric ground others optimizing one or another social objective. These solutions that we will critically review have more frequently been looked at on computational aspects. Our approach is at a general level that establishes at the outset an equivalence between an ordinal condition and additive optimization on the lattice of stable matchings. This equivalence charts out a domain where the correlates of ordinal criteria selected for equitability or social welfare appear as weights in optimization and vice versa. We call *convexity* the

ordinal condition in the equivalence and call the domain *modular*. Convexity involves preferences and requires that stable “mixtures” of matchings in a solution to also be in the solution. The modular domain is fully wide, separable across agents, and allows for computationally efficient optimization.

To introduce the framework, consider a society consisting of equal numbers of *men* and *women*. A matching (marriage) *problem* is a preference profile in which each agent has preferences over the opposite party. A *matching* uniquely assigns each man to a woman. Our primitive objects are *matching rules* that associate a nonempty set of matchings with each problem. The set of associated matchings can be thought of as a shortlist from which a final choice is to be made, the matchings that are assigned positive probability in a lottery, or the matchings used in a rotation scheme that, for example, specifies the periodical job assignments to the employees throughout their employment.

The central robustness criterion for a matching is *stability*, which requires that there is no unmatched man-woman pair who prefer each other to their assigned mates. In addition to requiring that a matching rule chooses only stable matchings, we impose *invariance under stability*, which requires the matchings chosen from two different problems be the same unless these problems induce different sets of stable matchings. We refer to these matching rules as *stable matching rules*.

Modularity of a stable matching rule requires the chosen matchings be the ones that optimize an explicit objective function, which is restricted to be *modular* by using the lattice structure of stable matchings for each problem. As demonstrated by several studies in the literature, formulating social objectives in the form a modular function provides analytical and computational tractability along with clarity and richness. Here, we provide several examples and results demonstrating that a wide range of objectives

can be implemented through these rules. As for analytical tractability, it follows from Proposition 4 together with the classical findings by Picard (1976) that optimizing a modular function over stable matchings boils down to finding the *minimum cuts* in a properly defined *flow network*. The latter problem has been extensively studied and known to be solvable efficiently.¹

The matching rules used in the literature can roughly be classified into two groups. In the first group, we can count the ones that choose the stable matchings optimizing an explicitly given objective function, which can be interpreted as a measure of social welfare or fairness—such as the utilitarian objective that seeks to maximize sum of individual utilities.² As discussed in Section 2.1, these rules typically violate *invariance under stability*, since the objective functions in question are formulated by using agents’ utilities/rankings over each other, independent of whether there is a stable matching in which these agents are matched or not. In the second group, we consider the rules whose formulations are not based on optimization of an explicitly given objective function, but formulated by directly using the mathematical structure of the stable matchings,³ and thus satisfy *invariance under stability*. Here, we provide a synthesis of these two groups of rules, in that modular stable matching rules have clear underlying objectives formulated in a tractable mathematical form, and satisfy *invariance under stability*.

¹See for example Picard & Queyranne (1980), Irving, Leather & Gusfield (1987), and Henzinger, Noe, Schulz & Strash (2020). More generally, Grötschel, Lovász & Schrijver (1981) show that minimizing a submodular function over a lattice family can be done in oracle-polynomial time via the ellipsoid algorithm.

²For other examples, one can count, *minimum regret stable matchings*, *egalitarian stable matchings*, the *minimum weight stable matchings*, *sex-equal stable matchings*, *rank maximal stable matchings*, and *balanced stable matchings*. For the related definitions we refer the reader to Manlove (2013, Chapter 1.3), Gusfield & Irving (1989), and the references therein.

³In Examples 1-3, we present three of such matching rules introduced by Teo & Sethuraman (1998), Cheng (2010) and Cheng, McDermid & Suzuki (2016).

In our analysis, we provide two characterizations for modular stable matching rules that reveal the ordinal content of optimizing a modular function under the constraint of *stability*. Thus, we provide testable conditions to verify if the observed choice of stable matchings in a society comply with optimization of a modular function. Additionally, these results guide us about normative qualitative criteria that one may consider to adopt a modular stable matching rule. On the technical side, we deepen and use the connection between the stable matchings lattice and the *rotations poset* (Irving & Leather 1986), which is used in computer science to design efficient matching algorithms, but seems to be overlooked in economics.

Our first axiom, *convexity*, requires that for each problem and for a given pair of matchings that are chosen by the rule, if one can form a stable matching by assigning all the agents to one of their mates in the given pair, then this newly formed stable matching should have been chosen as well. To introduce the second axiom, suppose that we transform a given problem—the preferences of the agents—such that all the agents move their mates in the matchings chosen by the rule to the top of their preferences by preserving the relative rankings elsewhere. Our *independence of irrelevant rankings* requires that if a matching that is stable in the original problem remains stable in this transformed problem, then it must be one of the matchings chosen by the rule in the initial problem.

In the second part of the paper, we propose a new equity notion that is based on the notion of the “median attainable mate”. To introduce this notion, consider an agent i and all the agents who are *attainable* for i in the sense that for each such agent there exists a stable matching in which i is assigned to him or her in the given problem. Next, from among the agents who are attainable for i , consider the one(s) with (a) *median*

rank.⁴ The median attainable mate for agent i is the (more preferred) attainable agent who has the lower attainable median rank. A matching rule satisfies *equity undominance* if for each problem, whenever a matching is chosen, there is no other stable matching in which each agent is assigned to a mate who is either the same or *closer* to agent's median attainable mate than compared to agent's mate at the chosen matching.

In Section 4, we present a simple problem in which there is a stable matching that assigns all the agents to their unique median attainable mates. However, several stable matching rules from the literature fail to choose this matching, and thus violate *equity undominance*. Then, in Theorem 2, we characterize the class of modular stable matching rules that satisfy *equity undominance*. The notion of *equity undominance* and Theorem 2 can be directly generalized by replacing median attainable mates by “ideal mates” as defined in Example 5. Our discussion in this part of the paper demonstrates that studying modular stable matching rule opens new avenues through which the issues of equity can be fruitfully analyzed. As for the many-to-one and many-to-many matchings, the set of stable matchings forms a distributive lattice under suitable restrictions.⁵ These findings together with the connection provided by Blair (1984)⁶ indicate that our results can be extended into more general matching contexts that are rich with market-design applications.

⁴If there is an even number of attainable agents for i , then there are two such agents.

⁵Alkan (2001) and Alkan & Gale (2003) show that in the context of many-to-one and many-to-many matchings, the distributivity of the stable matchings lattice is guaranteed by strengthening the *substitutability* of the choice functions with *size monotonicity* (called *law of aggregate demand* by Hatfield & Milgrom (2005)). Faenza & Zhang (2022) study algorithms for optimizing a modular function over the set of stable matchings in these models where preferences are replaced with choice functions.

⁶Who shows that for every distributive lattice, there is a marriage problem with a stable matchings lattice that is order-isomorphic to the given distributive lattice.

2 Modular Stable Matching rules

2.1 Stable matching rules

Let M be a set of n men and W be a set of n women. Each $m \in M$ has preferences over W and each $w \in W$ has preferences over M . Let $N = M \cup W$, then preferences of each agent $i \in N$ is represented by a **strict ordering**, which is denoted by \succ_i , i.e. \succ_i is *complete*, *transitive*, and *asymmetric*. Let \mathcal{P}_i denote the set of all possible preference relations for agent i , and $\mathcal{P} = \times_{i \in N} \mathcal{P}_i$ denote the set of all possible preference profiles. We denote a generic preference profile by \succ . A **matching** is a one-to-one function $\mu : M \cup W \rightarrow M \cup W$ such that for each $(m, w) \in M \times W$, we have $\mu(m) \in W$, $\mu(w) \in M$, and $\mu(m) = w$ if and only if $\mu(w) = m$.

A matching μ is **stable** at a problem $\succ \in \mathcal{P}$ if there is no **blocking-pair** $(m, w) \in M \times W$ such that $m \succ_w \mu(w)$ and $w \succ_m \mu(m)$. Let $\mathcal{S}(\succ)$ denote the set of all stable matchings at a given preference profile (problem) $\succ \in \mathcal{P}$. We often use \mathcal{S} instead of $\mathcal{S}(\succ)$ if the problem that is referred to is clear from the context. Let \triangleright_M denote the **men-wise better than** relation over \mathcal{S} , which are defined as follows: For each distinct $\mu, \mu' \in \mathcal{S}$, $\mu \triangleright_M \mu'$ if and only if for each $m \in M$, $\mu(m) \succ_m \mu'(m)$ or $\mu(m) = \mu'(m)$. The **women-wise better than** relation \triangleright_W is defined similarly. A **matching rule** is a mapping π that associates each problem $\succ \in \mathcal{P}$ with a nonempty set of matchings $\pi(\succ)$.

Definition. A **stable matching rule** π is a matching rule that satisfies the following two conditions:

Stability: For each problem $\succ \in \mathcal{P}$, $\pi(\succ) \subset \mathcal{S}(\succ)$.

Invariance under stability: For each $\succ, \succ' \in \mathcal{P}$, if $\mathcal{S}(\succ) = \mathcal{S}(\succ')$ then $\pi(\succ) = \pi(\succ')$.

Stability requires the matchings chosen for each problem be stable. *Invariance under stability* requires the matchings chosen from two different problems be the same unless these problems induce different set of stable matchings. Put differently, for a stable matching rule the set of stable matchings associated with each problem provides the relevant information to operate. Our Examples 4-7 demonstrate that to formulate an objective function whose optimization leads to a stable matching rule that satisfies *invariance under stability*, the critical notion is “attainable mates”. That is, a man and a woman are *attainable* for each other in a given problem, if there exists a stable matching in which they are matched to each other.

As noted in the introduction, several matching rules in the literature are formulated as to choose the stable matchings that optimize an objective function, which can be interpreted as a measure of social welfare or fairness. As a common theme in all of these rules, the objective function to be optimized is formulated via agents’ utilities/rankings over each other, independent of them being *attainable* or not. Therefore, they all fail to satisfy *invariance under stability*. To see this, let \succ be a problem and consider the problem \succ' that is obtained from \succ such that agents’ *attainable mates* are moved to top of their preferences by preserving the relative rankings elsewhere. It is easy to verify that the associated set of stable matchings remains the same. However, since \succ and \succ' can be distinct, optimization of an objective function based on agents’ utilities over each other for \succ and \succ' might result in different stable matchings. As demonstrated for the sex-equal stable matchings in Example 7, to remedy violation of *invariance under stability*, one needs to replace the use of agents’ rankings with their “attainable rankings”.

There is a another class of matching rules in the literature whose formulations are not explicitly based on optimization of an objective function, but are built on the

exquisite geometric structure of stable matchings. That is, for each pair of stable matchings μ and μ' , consider $\mu \vee \mu'$ ($\mu \wedge \mu'$) that maps each man to his best (worst) mate among the women he is matched to at μ or μ' ; it turns out that both $\mu \vee \mu'$ and $\mu \wedge \mu'$ are stable matchings as well, which in particular implies that the pair $\langle \mathcal{S}, \triangleright_M \rangle$ forms a **distributive lattice**.⁷ Next, we present three examples whose formulations are based on the structure of the stable matching lattice, satisfy invariance *under stability*, and thus form stable matching rules.

Example 1 (Median stable matchings). Let \succ be a problem with K stable matchings. For each man m , arrange his mates from these K stable matchings from his most preferred mate to his least preferred one. Let $w^k(m)$ denote the k -th woman in this sorted list, where the same woman can be counted multiple times. For each $k \in \{1, \dots, K\}$, define $\mu^k : M \cup W \rightarrow M \cup W$ such that $\mu^k(m) = w^k(m)$ for each $m \in M$. [Teo & Sethuraman \(1998\)](#) show that μ^k is a stable matching. Then, they define the *median stable matching(s)* as $\mu^{(K+1)/2}$ when K is odd, and $\mu^{K/2}$ and $\mu^{(K/2)+1}$ when K is even.⁸

Example 2 (Median of the stable matchings lattice). Consider the undirected graph associated with the stable matching lattice $\langle \mathcal{S}, \triangleright_M \rangle$, in which each $\mu \in \mathcal{S}$ is a vertex and for each $\mu, \mu' \in \mathcal{S}$, μ and μ' are *adjacent* if there is no $\mu'' \in \mathcal{S}$ with $\mu \triangleright_M \mu'' \triangleright_M \mu'$.⁹ For each $\mu, \mu' \in \mathcal{S}$, the *distance* between μ and μ' is the length of (the number of edges on) the shortest path (a.k.a. *geodesic*) between μ and μ' in this graph. [Cheng \(2010\)](#) analyzes stable matchings that are *median(s) of the stable matchings lattice* whose total or average distance from all other stable matchings is the least. She shows that when n is odd, the

⁷[Knuth \(1976\)](#), pp. 92-93, attributes the discovery of this lattice structure to J. H. Conway.

⁸The existence of (generalized) median stable matchings has been studied in other settings including: one-to-one matching with wages ([Schwarz & Yenmez 2011](#)), the college admissions model with responsive preferences ([Klaus & Klijn 2006a](#), [Sethuraman, Teo & Qian 2006](#)), the roommates problem ([Klaus & Klijn 2010](#)), many-to-many matching markets with contracts ([Chen, Egedal, Pycia & Yenmez 2016](#)).

⁹This graph is called the (undirected) Hasse diagram of $\langle \mathcal{S}, \triangleright_M \rangle$.

median stable matching is the unique median of the stable matchings lattice and when n is even, a stable matching μ is a median of the stable matchings lattice if and only if μ is between the median stable matchings according to \triangleright_M , i.e. $\mu^{(K/2)+1} \triangleright_M \mu \triangleright_M \mu^{K/2}$.

Example 3 (Center stable matchings). [Cheng, McDermid & Suzuki \(2016\)](#) formulate *center stable matching(s)* as the one(s) whose maximum distance (as described in Example 2) from any other stable matchings is the least. They provide a characterization of all center stable matchings, and show that a (specific) center stable matching can be computed in polynomial time.

2.2 Modularity

The main structural restriction that we would like to impose on a stable matching rule is *modularity*. Modularity requires optimization of an explicit objective function that is restricted in a specific way via the lattice structure of stable matchings. That is, for a given problem \succ , let $F : \mathcal{S}(\succ) \rightarrow \mathbb{R}$ be an **assessment function** that attaches a value $F(\mu)$ to each stable matching $\mu \in \mathcal{S}(\succ)$. Then, F is **modular** if for each $\mu, \mu' \in \mathcal{S}(\succ)$,

$$F(\mu) + F(\mu') = F(\mu \vee \mu') + F(\mu \wedge \mu'). \quad (1)$$

Note that (1) can be rewritten as $F(\mu) - F(\mu \wedge \mu') = F(\mu \vee \mu') - F(\mu')$. Then, the simple intuition behind modularity is as follows. The change from $\mu \wedge \mu'$ to μ corresponds to a group of men being matched to better women; the effect of this change should be the same if the change was made while another group of men were matched to better women (at μ') compared to their mates at $\mu \wedge \mu'$.

Definition. Let π be a stable matching rule. Then, π is **modular** if for each problem \succ , there exists a modular $F : \mathcal{S}(\succ) \rightarrow \mathbb{R}$ such that $\pi(\succ)$ is the set of stable matchings that minimize F , that is $\pi(\succ) = \operatorname{argmin}_{\mu \in \mathcal{S}(\succ)} F(\mu)$.

As another salient class of assessment functions, we can consider the ones that can be represented in additively separable form $\sum_{i \in N} F_i(\mu(i))$. This representation renders the direct interpretation that the social value of a stable matching is obtained by adding $F_i(\mu(i))$ for each agent i , which assesses the social value of matching agent i with $\mu(i)$. For example, imagine that a matching determines the partnership relation which can be between a senior and junior employee, or two teams running a joint project at the intersection of their areas of expertise. Then, the social value of matching agent i with $\mu(i)$ might be determined by the productivity of agent i when matched with $\mu(i)$. This formulation disallows complementarities, in the sense that the social value of matching agent i with $\mu(i)$ is the same regardless of other agents' matches. On the other hand, it follows from the findings of [Picard \(1976\)](#) and [Irving, Leather & Gusfield \(1987\)](#) that a stable matching optimizing such a given assessment function can be computed efficiently. In Proposition 1, we show that an assessment function is modular if and only if it can be represented as the sum of agents' *individual assessment functions* defined for each agent over his/her set of attainable mates. Formally, let \succ be a given problem, then a man m and a woman w are **attainable** for each other if $\mu(m) = w$ for some $\mu \in \mathcal{S}(\succ)$. For each $i \in N$, let A_i denote the set of attainable agents for i .

Proposition 1. Let $\succ \in \mathcal{P}$ be a problem and $F : \mathcal{S}(\succ) \rightarrow \mathbb{R}$ be an assessment function. Then, F is modular if and only if for each $i \in N$, there exists $F_i : A_i \rightarrow \mathbb{R}$ such that $F(\mu) = \sum_{i \in N} F_i(\mu(i))$ for each $\mu \in \mathcal{S}(\succ)$.

Here, we prove the if part of the statement. Suppose that for each $\mu \in \mathcal{S}(\succ)$, we have $F(\mu) = \sum_{i \in N} F_i(\mu(i))$, where $F_i : A_i \rightarrow \mathbb{R}$. To see that π is modular, let $i \in N$. Since for each $\mu, \mu' \in \mathcal{S}(\succ)$, we have $\{(\mu \vee \mu')(i), (\mu \wedge \mu')(i)\} = \{\mu(i), \mu'(i)\}$, we obtain that $\{F_i((\mu \vee \mu')(i)), F_i((\mu \wedge \mu')(i))\} = \{F_i(\mu(i)), F_i(\mu'(i))\}$. It directly follows that F_i is modular for each agent i , and thus F is modular since it is the sum of these modular functions. The proof of the only if part, which uses the *rotations poset*, is presented as Lemma 4 in Section 6.1.

Next, we present several examples to demonstrate the relevance, generality, and the possible limitations of the modular stable matching rules. The rules in these examples are based on aggregating agents' **attainable rankings**. That is, for each m and w who are attainable for each other, $Rank_m^A(w)$ ($Rank_w^A(m)$) is the rank of w (m) in $\succ_m \mid_{A_m}$ ($\succ_w \mid_{A_w}$), which is obtained by restricting \succ_m (\succ_w) to the women (men) who are attainable for m (w).

Example 4 (Maximizing total attainable ranks). In the vein of utilitarian welfare measures, it may be reasonable to evaluate each stable matching according to the sum of agents' attainable ranks in the matching. That is, for each problem \succ , let $\pi(\succ)$ be the set of stable matchings that maximize $\sum_{mw \in \mu} (Rank_m^A(w) + Rank_w^A(m))$. In Lemma 1 of Section 6.1, we show that this sum is constant among all stable matchings, and therefore does not differentiate any stable matching from the others.

Example 5 (Minimizing total spread from the ideals). For each agent $i \in N$, from among the agents who are attainable for i , let $I(i)$ be the *ideal* partner for i in the sense that assigning i to $I(i)$ makes agent i reach the welfare level that is found ideal for him. We allow two different agents have the same ideal partner. For a given stable matching μ , we can measure the **spread from the ideal** for agent i by $|Rank_i^A(\mu(i)) - Rank_i^A(I(i))|$.

Then, consider the stable matching rule π choosing the set of stable matchings that minimize the *total spread from the ideals*. That is, for each problem \succ , let $\pi(\succ)$ be the set of matchings that minimizes $\sum_{i \in N} |\text{Rank}_i^A(\mu(i)) - \text{Rank}_i^A(I(i))|$. It directly follows from Proposition 1 that π is modular.

Example 6 (Minimizing total spread from attainable medians). We may add further structure into the previous example as follows. For each agent i , from among the agents who are attainable for i , consider the one(s) with (a) *median rank*. Note that if there is an odd number of attainable agents for i , then there is a unique attainable agent with this property, otherwise there are two such agents. Let med_i^A be the (more preferred) attainable agent with the lowest attainable median rank, i.e. $\text{Rank}_i^A(\text{med}_i^A) = \lfloor |A_i|/2 \rfloor$. Suppose that med_i^A is viewed as the ideal partner for i . Then, consider the stable matching rule π choosing the set of stable matchings that minimize the total spread from the median. That is, for each problem \succ , let $\pi(\succ)$ be the set of matchings that minimizes $\sum_{i \in N} |\text{Rank}_i^A(\mu(i)) - \text{Rank}_i^A(\text{med}_i^A)|$. Alternatively, one can also consider the stable matching rule choosing the set of stable matchings that maximize the total spread from the median. It is easy to verify that in this rule the extremal matchings are chosen at every problem, possibly together with other stable matchings. It follows from [Gusfield \(1987\)](#)¹⁰ that identifying the set of attainable mates is a polynomial task. Then, by using the findings of [Irving, Leather & Gusfield \(1987\)](#), one can show that a stable matching minimizing the total spread from median can be computed in polynomial time, even in the absence of the *oracle*.¹¹

¹⁰[Gusfield \(1987\)](#) shows that a representation for the rotation poset, $\langle \mathcal{R}, \rightarrow \rangle$ as denoted in Section 6.1, can be constructed in $O(n^2)$ time. Our discussion in Section 6.1 will clarify the connection between attainable mates and the rotation poset.

¹¹It follows from [Irving, Leather & Gusfield \(1987\)](#) [Theorem 5.2] that optimizing an additive function over the rotation poset can be computed in $O(n^2)$ time. Since, our Lemma 2 in Section 6.1 shows that a modular assessment function can be represented as an additive function over the rotation poset, the conclusion follows.

Example 7 (Minimizing the difference between total attainable ranks). As a counterpart of the *sex-equal stable matchings* (Gusfield & Irving 1989), consider the stable matching rule π choosing the set of *attainable sex-equal stable matchings* that minimize the absolute value of the difference between each sides' total attainable ranks. That is, for each problem \succ , let $\pi(\succ)$ be the set of matchings that minimizes $|\sum_{m \in M} \text{Rank}_m^A(\mu(m)) - \sum_{w \in W} \text{Rank}_w^A(\mu(w))|$. We will show that this stable matching rule is not modular.

3 Axioms and characterizations

3.1 Convexity

We present two characterizations for modular stable matching rules revealing the ordinal content of optimizing a modular function. Our first axiom is *convexity*, which requires that for a given pair of stable matchings that are chosen by the rule at a problem, if one can form a “mixture” stable matching by assigning agents to one of their mates in the given pair, then this newly formed matching should be chosen as well.

Convexity: For each problem $\succ \in \mathcal{P}$, if $\mu', \mu'' \in \pi(\succ)$ and there exists $\mu \in \mathcal{S}(\succ)$ such that $\mu(m) \in \{\mu'(m), \mu''(m)\}$ for each $m \in M$, then $\mu \in \pi(\succ)$.

Note that *convexity* is weaker than requiring that all the stable matchings that are between the chosen matchings—according to the men-wise better than relation—must be chosen as well, which is also referred to as convexity in lattice theory literature. To see this, consider a stable matching rule π such that for each problem \succ , only the extremal matchings are chosen whenever there is no other $\mu \in \mathcal{S}(\succ)$ such that $\mu(m) \in \{\mu^M(m), \mu^W(m)\}$ for each $m \in M$ (such as the problem in Example 8); and chooses all

of the stable matchings, otherwise. Although π satisfies our *convexity*, it clearly violates the latter requirement. It is worth to emphasize that *convexity* requires a matching that is a mixture of two chosen stable matchings be chosen only if this mixture matching is also stable.¹² In Section 8.1, in order to understand the notion of mixtures and stable mixtures better, we make two structural observations. In Lemma 12, we provide a simple procedure to obtain all the mixtures of two stable matchings. Then, in Lemma 13, we show that the set of stable mixtures of two stable matchings is a *Boolean* sublattice of the stable matchings, which can be interpreted as having a symmetric structure.

Theorem 1. *Let π be a stable matching rule. Then, π is modular if and only if π satisfies convexity.*

Proof. Please see Section 7.1. □

Proof sketch for the if part: Let π be a stable matching rule that satisfies *convexity*. To simplify the construction of the desired modular assessment function F , we use the connection between the stable matchings and the so-called *closed* sets of *rotations*. Rotations—first introduced by Irving (1985)—can be intuitively thought of as the incremental changes that transform a stable matching μ into another stable matching μ' such that there lies no other stable matching between the two (according to the men-wise domination relation). A rotation ρ *exposed in* a stable matching μ is an (ordered) cyclic sequence of distinct man-woman pairs $[(m_1, w_1), (m_2, w_2), \dots, (m_k, w_k)]$ such that each m_i is matched to w_i in μ . To *eliminate* ρ and obtain μ' , each man m_i in ρ is matched to w_{i+1} . A rotation ρ *precedes* another rotation ρ' , if ρ must be eliminated first in order to obtain a stable matching in which ρ' is exposed. A set of rotations R is *closed* if whenever

¹² The example in Section 8.2 demonstrates that every mixture of stable matchings is not necessarily stable.

a rotation ρ is contained in R , then all the rotations that precede ρ are also contained in R . In their main result, [Irving & Leather \(1986\)](#) show that each stable matching μ is associated with a unique closed set of rotations R_μ . It follows from this result and few simple observations that we made in Section 6.1 that constructing the desired F is equivalent to assigning a weight $g(\rho)$ to each rotation ρ such that $\pi(\succ)$ is the set of stable matchings that minimize $\sum_{\rho \in R_\mu} g(\rho)$.




Figure 1: A demonstration of our weight assignment to the rotations, where members of $\pi(\succ)$ are lightly colored.

As the second step of the proof, we introduce *hyper-rotations*, generically denoted by λ , which can be thought of as sets of rotations connecting the chosen stable matchings (see Section 6.2). Since, by *convexity*, $\pi(\succ)$ is a sublattice, let $\bar{\mu}(\underline{\mu})$ be its \triangleright_M -best(worst) matching. Now, there can be other chosen stable matchings between $\bar{\mu}$ and $\underline{\mu}$ that are obtained by eliminating hyper-rotations. Therefore, the weights of the rotations should be assigned such that the total weight of each hyper-rotation is zero, while the total weight of each (relatively) closed set of rotations $R \subsetneq \lambda$ —which correspond to the unchosen

stable matchings between $\bar{\mu}$ and $\underline{\mu}$ is positive. For a fixed hyper-rotation λ , achieving this requires “preloading” in the sense that we assign positive weights to the rotations with no predecessors in λ and assign negative weights (say -1) to the rotations with no successors in λ , while assigning zero weights to all other rotations in λ .

In Figure 1, we demonstrate this construction for a particularly simple case. For the general construction, the inner structure of hyper-rotations, and therefore *convexity*, plays a crucial role. In Lemma 8, we show that *convexity* implies that each hyper-rotation λ is a *connected* set of rotations. This observation paves the way to preload the positive weights in a hyper-rotation properly, in that we can distribute each negative weight assigned to a rotation ρ with no successors equally among the rotations that have no predecessors and precede ρ . We complete the proof by showing that we obtain the desired weight assignments through this construction.

Since Ore (1956),¹³ it is known that any finite distributive lattice can be obtained as the minimizers of a submodular function¹⁴ that is defined over the subsets of a finite set.¹⁵ This raises the question that if we can provide a counterpart to Theorem 1 for stable matching rules that are *submodular*. That is, the chosen stable matchings for each problem \succ , are the ones that minimize a **submodular** assessment function F : for each $\mu, \mu' \in \mathcal{S}(\succ)$, we have $F(\mu) + F(\mu') \geq F(\mu \vee \mu') + F(\mu \wedge \mu')$. The following corollary to Theorem 1 shows that the answer is in the affirmative.

Corollary 1. *Let π be a stable matching rule. Then, π is submodular if and only if $\pi(\succ)$ is a sublattice of $\mathcal{S}(\succ)$ for each problem $\succ \in \mathcal{P}$.*

¹³Also see Lemma 2.1 in Fujishige (2005).

¹⁴For an introduction to submodular functions, one can consult Lovász (1983). Edmonds (1970) and Topkis (1978) are two classical references on the minimization of submodular functions.

¹⁵In a recent paper, Fujii & Kijima (2019) sharpens this result by showing that any finite distributive lattice appears as the minimizers of a M^\sharp -concave (Murota 1998) set function. It is not immediately clear how to formulate M^\sharp -concavity for an assessment function, which is defined over stable matchings.

Proof. The only if part directly follows from π being submodular. Conversely, let \succ be a problem, and consider $\pi(\succ)$. Since $\pi(\succ)$ is a sublattice of $\mathcal{S}(\succ)$, let $\bar{\mu}(\underline{\mu})$ be the \triangleright_M -best(worst) matching in $\pi(\succ)$. Then, define $L(\succ) = \{\mu \in \mathcal{S}(\succ) \mid \bar{\mu} \triangleright_M \mu \triangleright_M \underline{\mu}\}$. Note that $L(\succ)$ is a *convex* sublattice of $\mathcal{S}(\succ)$ that contains $\pi(\succ)$. Then, it follows from Theorem 1 that there exists a modular assessment function F such that $L(\succ)$ is the set of matchings that minimizes F . Next, define a new assessment function F^* such that for each $\mu \in \mathcal{S}(\succ)$, if $\mu \in \pi(\succ)$, then $F^*(\mu) = F(\mu) - \epsilon$ for some fixed $\epsilon > 0$; and $F^*(\mu) = F(\mu)$, otherwise. It directly follows from this construction that $\pi(\succ)$ is the set of stable matchings that minimize F^* .

To see that F^* is submodular, note that $F^*(\mu) + F^*(\mu') < F^*(\mu \vee \mu') + F^*(\mu \wedge \mu')$ for some $\mu, \mu' \in \mathcal{S}(\succ)$ only if $\mu \in \pi(\succ)$ and $\mu' \notin \pi(\succ)$ (or vice versa). But, then it follows from a basic observation—Lemma 5 in Section 6.1—that $\mu \vee \mu' \in \pi(\succ)$ or $\mu \wedge \mu' \in \pi(\succ)$, indicating that this is not possible. \square

3.2 Independence of irrelevant rankings

To introduce our second axiom, we need the notion of a π -transformed problem. For each problem $\succ \in \mathcal{P}$ and each agent $i \in N$, let $\pi_i(\succ)$ be the set of agents that i is assigned to at any matching $\mu \in \pi(\succ)$, i.e. $\pi_i(\succ) = \{\mu(i) \in N \mid \mu \in \pi(\succ)\}$. Then, the **π -transformed problem** \succ^π is the problem obtained from \succ such that for each agent $i \in N$, each member of $\pi_i(\succ)$ is moved to the top of agent i 's preferences by preserving the relative rankings elsewhere.

Now, let us make a simple observation. For each $\mu \in \mathcal{S}(\succ)$, if $\mu \in \pi(\succ)$, then in transforming \succ into \succ^π , for each $i \in N$, the set of agents that i prefers to $\mu(i)$ remains

the same or shrinks, i.e. $\{j \mid j \succ_i^\pi \mu(i)\} \subset \{j \mid j \succ_i \mu(i)\}$. Therefore, for each $\mu \in \mathcal{S}(\succ)$, if $\mu \in \pi(\succ)$ then $\mu \in \mathcal{S}(\succ^\pi)$. Our next axiom requires the converse, that is, if a stable matching remains stable after the transformation, then it must be one of the matchings chosen by the rule in the initial problem. It follows from Proposition 2 that *convexity* and *independence of irrelevant rankings* both independently characterize modular stable matching rules.

Independence of Irrelevant Rankings (IIR): For each problem $\succ \in \mathcal{P}$ and $\mu \in \mathcal{S}(\succ)$, if $\mu \in \mathcal{S}(\succ^\pi)$, then $\mu \in \pi(\succ)$.

Proposition 2. *Let π be a stable matching rule. Then, π satisfies convexity if and only if π satisfies independence of irrelevant rankings.*

Proof. Please see Section 7.2. □

To see one side of the connection between *convexity* and *IIR*, we show that if a stable matching rule π satisfies *IIR*, then π satisfies *convexity*. For a given problem \succ , let $\mu', \mu'' \in \pi(\succ)$ and $\mu \in \mathcal{S}(\succ)$ such that for each $m \in M$, $\mu(m) \in \{\mu'(m), \mu''(m)\}$. Since π satisfies *IIR*, we have $\pi(\succ) = \mathcal{S}(\succ) \cap \mathcal{S}(\succ^\pi)$. Therefore, to conclude that $\mu \in \pi(\succ)$ it is sufficient to show that $\mu \in \mathcal{S}(\succ^\pi)$. By contradiction, suppose that there is a blocking pair (m, w) . Therefore, $w \succ_m^\pi \mu(m)$ and $m \succ_w^\pi \mu(w)$. Now, since we have $\mu(m) \in \{\mu'(m), \mu''(m)\}$ and $\mu(w) \in \{\mu'(w), \mu''(w)\}$, where $\mu', \mu'' \in \pi(\succ)$, by the definition of the π -transformation, no woman is moved over $\mu(m)$ in m 's preferences and no man is moved over $\mu(w)$ in w 's preferences while moving from \succ into \succ^π . Then, it follows from $w \succ_m^\pi \mu(m)$ and $m \succ_w^\pi \mu(w)$ that $w \succ_m \mu(m)$ and $m \succ_w \mu(w)$, contradicting that $\mu \in \mathcal{S}(\succ)$.

We conclude this section by revisiting the stable matching rules from the literature that we discussed in Section 2.1. By using our Theorem 1, we can easily check if these

M \ W	w_1	w_2	w_3	w_4
m_1	1 4	4 1	2 3	3 2
m_2	4 1	1 4	3 2	2 3
m_3	3 2	2 3	1 4	4 1
m_4	2 3	3 2	4 1	1 4




Figure 2: The problem and the associated stable matchings lattice.

rules are modular or not. It turns out that the matching rule that chooses the median(s) of the stable matching lattice (Example 2) is modular since it is *convex*. To see this, recall that that this rule either chooses the unique median stable matching or all the stable matchings that are between the median stable matchings according to the men-wise better than relation. However, the following simple example demonstrates that the rules presented in Examples 1,3, and 7 fail to satisfy *convexity*.

Example 8. Consider the problem with eight agents whose preferences are represented by the table in Figure 2 such that each entry ij is associated with man m_i and woman w_j , the \succ_{m_i} -rank of w_j is written in the bottom corner and the rank of m_i in \succ_{w_j} is written in the top corner. in Figure 2, we also present the associated stable matchings lattice such that each stable matching is represented as an array $[r_1, \dots, r_4]$, where each r_i is the \succ_{m_i} -rank of the women who is matched with m_i . Then, $[2222]$ and $[3333]$ are the median stable matchings, and the attainable sex-equal matchings. The center of the stable matching lattice are $[3322]$ and $[2233]$. Thus, all three rules fail to satisfy *convexity*.

4 A new (class of) equity notions

We believe that studying modular stable matching rule opens new avenues through which the issues of equity can be fruitfully analyzed. To demonstrate this, let us consider the problem and its stable matchings lattice presented in Figure 3. We follow the notation used in Example 8 with the addition that if a pair is unattainable, then the associated entry is shadowed.




Figure 3: The problem and the associated stable matchings lattice.

It is easy to see that the unique median stable matching, the unique median of the stable matching lattice, and the unique center stable matching is the one in which each man is matched to his second-ranked woman and each woman is matched to her fifth-ranked man. However, one can argue that the matching in which each agent–man or woman–is matched to his/her third-ranked attainable mate is more equitable, since each agent is matched to his/her median attainable mate.

To formalize this intuition as a principle, let i be an agent and $j, j' \in A_i$ be a pair of attainable mates for i . Then, agent j is **closer to** med_i^A than agent j' if $|Rank_i^A(j) -$

$|Rank_i^A(med_i^A)| < |Rank_i^A(j') - Rank_i^A(med_i^A)|$. Our next axiom requires that if a matching rule chooses a stable matching μ , then there should be no other stable matching in which each agent with a different mate is assigned to someone who is closer to his/her median attainable mate than compared to his/her mate at μ .

Equity undominance: For each problem $\succ \in \mathcal{P}$, if $\mu \in \pi(\succ)$, then there is no $\mu' \in \mathcal{S}(\succ)$ such that for each $i \in N$ with $\mu(i) \neq \mu'(i)$ we have $\mu'(i)$ is closer to med_i^A than $\mu(i)$.

In our previous example, this principle uniquely pins down the matching in which each agent is matched to his/her third-ranked mate. Next, we characterize the class of modular stable matching rules that satisfy *equity undominance*. Let i be an agent, then the individual assessment function $F_i : A_i \rightarrow \mathbb{R}$ is **unimodal** with mode med_i^A if F_i is monotonically increasing for $med_i^A \succ_i j$ and monotonically decreasing for $j \succ_i med_i^A$.¹⁶

Theorem 2. A modular stable matching rule π satisfies equity undominance if and only if for each problem $\succ \in \mathcal{P}$, $\pi(\succ)$ is the set of stable matchings that minimize $-\sum_{i \in N} F_i(\mu(i))$, where $F_i : A_i \rightarrow \mathbb{R}$ is unimodal with mode med_i^A for each $i \in N$.

Proof. Please see Section 7.3. □

As a stable matching rule, one can consider to choose all the equity undominated stable matchings for each problem. However, as demonstrated in Section 8.2, the set of equity undominated stable matchings turns out to be rather unstructured, in particular, it fails to satisfy *convexity*.¹⁷ As for another stable matching rule that fails to satisfy *equity undominance*, we can consider the stable matching rule presented in Example 7,

¹⁶Put differently, F_i attains its maximum at med_i^A , and for each $j, j' \in A_i \setminus \{med_i^A\}$, we have $F_i(j) > F_i(j')$ if j' is further away from med_i^A compared to j according to \succ_i , i.e. $j' \succ_i j \succ_i med_i^A$ or $med_i^A \succ_i j \succ_i j'$.

¹⁷Additionally, this set is not a sublattice of the original problem in general, and a stable matching that is between two equity undominated matching, according to the men-wise better than relation, can be equity dominated. These points are demonstrated via the problem presented in Section 8.2.

which chooses the set of attainable sex-equal stable matchings.¹⁸ On the other hand, the objective function of total spread from attainable medians formulated in Example 6, can be expressed as a member of this additive unimodal family, where for each agent i and stable matching μ , $F_i(\mu(i)) = -|Rank_i^A(\mu(i)) - Rank_i^A(med_i^A)|$.

Remark 1. The notion of *equity undominance* and Theorem 2 can be generalized by replacing median attainable mates by ideal mates as defined in Example 5. Thus, we obtain a rich class of equity notions in which ideal mates are freely specified. On the other hand, requiring that each agent is attached to a unique ideal mate can be a rather demanding feature that rules out reasonable modular stable matching rules such as the one presented in Example 2.

5 Related literature and final comments

We proposed a general foundational framework to address the issues of social welfare and equity in the stable matching model and presented axiomatic characterizations for modular stable matching rules. Existing studies have focused only on rules that singles out an extremal matching, including Balinski & Sönmez (1999), Ehlers & Klaus (2006) and Kojima & Manea (2010) who offer different characterizations of the Gale-Shapley student-optimal stable matching rule. A general axiomatic approach to the problem of fair algorithms was presented by Masarani & Gokturk (1989). However, their approach concentrates on the algorithm, not on the resulting matchings, and concludes with an impossibility result. Relatedly, Klaus & Klijn (2006b) introduce the *procedural fairness* notion that labels a probabilistic stable matching rules as fair if each agent has the

¹⁸See the problem presented in Section 8.2, where the unique attainable sex-equal stable matching is equity dominated.

same probability to move at a certain point in the procedure that determines the final probability distribution. Their approach does not provide any criterion for fairness or equitability of a stable matching.

Modularity properties of agents' preferences was investigated by [Kreps \(1979\)](#) in decision-theory literature and by [Milgrom & Shannon \(1994\)](#) in the monotone comparative statics literature. [Chambers & Echenique \(2008\)](#) clarifies the connection between these two different approaches. The closest papers to ours are [Kreps \(1979\)](#) and [Chambers & Echenique \(2009\)](#) who provide representations for modular preferences over lattices under the additional assumption of *monotonicity*. However, violation of *monotonicity* is the departure point of our question, since a stable matching rule that satisfy *monotonicity* would choose one of the extremal matchings. As demonstrated in our proofs, here, the neat mathematical structure of the stable matchings paves the way for our results in the absence of *monotonicity*. As another key modelling difference, our primitives are not agents' preferences but stable matching rules that can be thought of as choice rules over the set of stable matchings.

In the second part of the paper, we analyze a novel equity notion to demonstrate that the class of modular stable matching rules are rich enough to formulate and analyze similar notions of equity. This new framework leads to a wide variety of open problems to be solved. To name a few, a major concern for stable matching rules is to provide the agents with the right incentives to report their preferences truthfully. Recall that the set of matchings chosen by a rule can be interpreted as the ones that are assigned positive probability in a lottery. Therefore, it seems reasonable to consider probabilistic assignment rules, as formulated by [Bogomolnaia & Moulin \(2001\)](#), associated with a modular stable matching rule and investigate general conditions that guarantee its

strategy-proofness in terms of stochastic-dominance. We have observed that there are modular stable matching rules, such as the one presented in Example 6, that can be implemented efficiently. In this vein, another direction that we find worth to explore is discovering the principles underlying the general class of efficiently implementable rules.

References

- Alkan, A. (2001), ‘On preferences over subsets and the lattice structure of stable matchings’, *Review of Economic Design* **6**(1), 99–111. [7](#)
- Alkan, A. & Gale, D. (2003), ‘Stable schedule matching under revealed preference’, *Journal of Economic Theory* **112**(2), 289–306. [7](#)
- Balinski, M. & Sönmez, T. (1999), ‘A tale of two mechanisms: Student placement’, *Journal of Economic Theory* **84**, 73–94. [24](#)
- Birkhoff, G. (1937), ‘Rings of sets’, *Duke Mathematical Journal* **3**(3), 443–454. [33](#)
- Blair, C. (1984), ‘Every finite distributive lattice is a set of stable matchings’, *Journal of Combinatorial Theory, Series A* **37**(3), 353–356. [7](#)
- Bogomolnaia, A. & Moulin, H. (2001), ‘A new solution to the random assignment problem’, *Journal of Economic theory* **100**(2), 295–328. [25](#)
- Chambers, C. P. & Echenique, F. (2008), ‘Ordinal notions of submodularity’, *Journal of Mathematical Economics* **44**(11), 1243–1245. [25](#)

- Chambers, C. P. & Echenique, F. (2009), ‘Supermodularity and preferences’, *Journal of Economic Theory* **144**(3), 1004–1014. [25](#)
- Chen, P., Egedal, M., Pycia, M. & Yenmez, M. B. (2016), ‘Median stable matchings in two-sided markets’, *Games and Economic Behavior* **97**, 64–69. [10](#)
- Cheng, C. T. (2010), ‘Understanding the generalized median stable matchings’, *Algorithmica* **58**(1), 34–51. [5](#), [10](#)
- Cheng, C. T., McDermid, E. & Suzuki, I. (2016), ‘Eccentricity, center and radius computations on the cover graphs of distributive lattices with applications to stable matchings’, *Discrete Applied Mathematics* **205**, 27–34. [5](#), [11](#)
- Edmonds, J. (1970), ‘Submodular functions, matroids, and certain polyhedra, combinatorial structures and their applications’, *New York* pp. 69–87. [18](#)
- Ehlers, L. & Klaus, B. (2006), ‘Efficient priority rules’, *Games and Economic Behavior* **55**(2), 372–384. [24](#)
- Faenza, Y. & Zhang, X. (2022), ‘Affinely representable lattices, stable matchings, and choice functions’, *Mathematical Programming* pp. 1–40. [7](#)
- Fujii, T. & Kijima, S. (2019), ‘Any finite distributive lattice is isomorphic to the minimizer set of an M^{\natural} -concave set function’, *arXiv preprint arXiv:1903.08343* . [18](#)
- Fujishige, S. (2005), *Submodular functions and optimization*, Elsevier. [18](#)
- Gale, D. & Shapley, L. S. (1962), ‘College admissions and the stability of marriage’, *American Mathematical Monthly* **69**, 9–15. [3](#)

- Grötschel, M., Lovász, L. & Schrijver, A. (1981), ‘The ellipsoid method and its consequences in combinatorial optimization’, *Combinatorica* **1**(2), 169–197. [5](#)
- Gusfield, D. (1987), ‘Three fast algorithms for four problems in stable marriage’, *SIAM Journal on Computing* **16**(1), 111–128. [14](#)
- Gusfield, D. & Irving, R. W. (1989), ‘The stable marriage problem: structure and algorithms’. [5](#), [15](#)
- Hatfield, J. W. & Milgrom, P. (2005), ‘Matching with contracts’, *American Economic Review* **95**, 913–935. [7](#)
- Henzinger, M., Noe, A., Schulz, C. & Strash, D. (2020), ‘Finding all global minimum cuts in practice’, *arXiv preprint arXiv:2002.06948* . [5](#)
- Irving, R. W. (1985), ‘An efficient algorithm for the stable roommates problem’, *Journal of Algorithms* **6**(4), 577–595. [16](#), [31](#)
- Irving, R. W. & Leather, P. (1986), ‘The complexity of counting stable marriages’, *SIAM Journal on Computing* **15**(3), 655–667. [6](#), [17](#), [33](#), [34](#)
- Irving, R. W., Leather, P. & Gusfield, D. (1987), ‘An efficient algorithm for the optimal stable marriage’, *Journal of the ACM (JACM)* **34**(3), 532–543. [5](#), [12](#), [14](#)
- Klaus, B. & Klijn, F. (2006a), ‘Median stable matching for college admissions’, *International Journal of Game Theory* **34**(1), 1–11. [10](#)
- Klaus, B. & Klijn, F. (2006b), ‘Procedurally fair and stable matching’, *Economic Theory* **27**(2), 431–447. [24](#)

- Klaus, B. & Klijn, F. (2010), ‘Smith and Rawls share a room: stability and medians’, *Social Choice and Welfare* **35**(4), 647–667. [10](#)
- Knuth, D. E. (1976), ‘Marriages stables’, *Technical report* . [10](#)
- Kojima, F. & Manea, M. (2010), ‘Axioms for deferred acceptance’, *Econometrica* **78**, 633–653. [24](#)
- Kreps, D. M. (1979), ‘A representation theorem for “preference for flexibility”’, *Econometrica: Journal of the Econometric Society* pp. 565–577. [25](#)
- Lovász, L. (1983), Submodular functions and convexity, in ‘Mathematical programming the state of the art’, Springer, pp. 235–257. [18](#)
- Manlove, D. (2013), *Algorithmics of matching under preferences*, Vol. 2, World Scientific. [5](#)
- Masarani, F. & Gokturk, S. S. (1989), ‘On the existence of fair matching algorithms’, *Theory and Decision* **26**(3), 305–322. [24](#)
- Milgrom, P. & Shannon, C. (1994), ‘Monotone comparative statics’, *Econometrica: Journal of the Econometric Society* pp. 157–180. [25](#)
- Murota, K. (1998), ‘Discrete convex analysis’, *Mathematical Programming* **83**(1), 313–371. [18](#)
- Ore, O. (1956), ‘Studies on directed graphs, I’, *Annals of Mathematics* pp. 383–406. [18](#)
- Picard, J.-C. (1976), ‘Maximal closure of a graph and applications to combinatorial problems’, *Management Science* **22**(11), 1268–1272. [5](#), [12](#)

- Picard, J.-C. & Queyranne, M. (1980), On the structure of all minimum cuts in a network and applications, in ‘Combinatorial Optimization II’, Springer, pp. 8–16. [5](#)
- Schwarz, M. & Yenmez, M. B. (2011), ‘Median stable matching for markets with wages’, *Journal of Economic Theory* **146**(2), 619. [10](#)
- Sethuraman, J., Teo, C.-P. & Qian, L. (2006), ‘Many-to-one stable matching: geometry and fairness’, *Mathematics of Operations Research* **31**(3), 581–596. [10](#)
- Teo, C.-P. & Sethuraman, J. (1998), ‘The geometry of fractional stable matchings and its applications’, *Mathematics of Operations Research* **23**(4), 874–891. [5](#), [10](#)
- Topkis, D. M. (1978), ‘Minimizing a submodular function on a lattice’, *Operations Research* **26**(2), 305–321. [18](#)

6 Appendix A: Stepping stones

6.1 Rotations and some preliminary observations

For a given fixed problem $\succ \in \mathcal{P}$, *rotations*—first introduced by [Irving \(1985\)](#)—are the incremental changes that need to be made so that a stable matching μ can be transformed into another stable matching μ' such that $\mu \triangleright_M \mu'$ and there is no other stable matching μ'' such that $\mu \triangleright_M \mu'' \triangleright_M \mu'$.

$$\begin{aligned} \rho^{11} &= [(m_1, w_1), (m_2, w_2)] \\ \rho^{12} &= [(m_3, w_3), (m_4, w_4)] \\ \rho^{13} &= [(m_5, w_5), (m_6, w_6)] \\ \rho^2 &= [(m_1, w_2), (m_4, w_3), (m_5, w_6), (m_2, w_1), (m_3, w_4), (m_6, w_5)] \\ \rho^3 &= [(m_1, w_3), (m_2, w_4), (m_3, w_5), (m_4, w_6), (m_5, w_1), (m_6, w_2)] \\ \rho^4 &= [(m_1, w_4), (m_2, w_5), (m_3, w_6), (m_4, w_1), (m_5, w_2), (m_6, w_3)] \end{aligned}$$

M \ W	w_1	w_2	w_3	w_4	w_5	w_6
m_1	1	6	2	5	3	4
m_2	2	5	1	6	2	4
m_3	5	1	6	2	3	4
m_4	4	3	5	1	6	2
m_5	3	4	2	5	1	6
m_6	6	2	4	3	5	1

Figure 4: The problem in Section 4 and the associated rotations.

Let μ^M and μ^W denote the men-optimal and women-optimal stable matchings. Let μ be a stable matching such that $\mu \neq \mu^W$. Then, $\mu(m) \neq \mu^W(m)$ for some man m . For each such man m , define his **successor woman** at μ , denoted by $s_\mu(m)$, as the first attainable woman for m on his preference list who is less preferred than $\mu(m)$ (and prefers him over her current partner in μ) i.e. $s_\mu(m)$ is the \succ_m -best attainable woman w such that $\mu(m) \succ_m w$ (and $m \succ_w \mu(m)$). A **rotation** ρ **exposed in** μ is an (ordered) cyclic sequence of distinct man-woman pairs $\rho = [(m_1, w_1), (m_2, w_2), \dots, (m_k, w_k)]$ such that $m_i w_i \in \mu$ and $s_\mu(m_i) = w_{i+1}$ for each $i \in \{1, \dots, k\}$, where the addition in the subscripts is modulo k . To **eliminate** a rotation ρ exposed in a stable matching μ , each man m_i in ρ is matched to w_{i+1} while all the pairs that are not in ρ are kept the same. As a result, we obtain another stable matching, denoted by $\mu \circ \rho$, such that $\mu \triangleright_M \mu \circ \rho$

and there is no other stable matching μ' with $\mu \triangleright_M \mu' \triangleright_M \mu \circlearrowright \rho$.

Let \mathcal{R} denote the set of all rotations. A rotation ρ **precedes** another rotation ρ' , denoted by $\rho \rightarrow \rho'$, if in order to obtain a stable matching in which ρ' is exposed, ρ must be eliminated first. We assume that a rotation precedes itself. A rotation ρ **immediately precedes** another rotation ρ' if $\rho \rightarrow \rho'$ and there is no other rotation ρ'' such that $\rho \rightarrow \rho'' \rightarrow \rho'$. A distinct pair of rotations ρ and ρ' are **independent** if none of them precedes the other. The pair $\langle \mathcal{R}, \rightarrow \rangle$ is called the **rotation poset**.




Figure 5: The stable matchings lattice and the rotation poset for the problem in Section 4 .

A subset of \mathcal{R} , generically denoted by R , is **closed** if whenever a rotation $\rho \in R$, then all the rotations that precede ρ are also in R . Let $Cl(\mathcal{R})$ denote the set of all closed subsets of \mathcal{R} . We suppress the reference to the specific problem $\succ \in \mathcal{P}$, whenever it is clear from the context.

We will use the following basic properties of rotations. First, a man-woman pair (m, w) belongs to a rotation if and only if it appears in some stable matching (i.e. if they are attainable) and w is not the worst mate of m in all stable matchings; and a man-woman pair can be an element of at most one rotation (Irving & Leather 1986).

For each $\rho \in \mathcal{R}$, let N_ρ denote the set of agents who appear in rotation ρ . We note that if a pair of rotations ρ and ρ' are independent, then there is no agent who appears both in ρ and ρ' , i.e. $N_\rho \cap N_{\rho'} = \emptyset$. Conversely, if there is no agent who appears both in ρ and ρ' , then none of these rotations immediately precedes the other.

In their main result, Irving & Leather (1986) show that the closed subsets of \mathcal{R} endowed with the set containment relation $\langle Cl(\mathcal{R}), \subset \rangle$ is a lattice that is *order isomorphic*¹⁹ to $\langle S, \triangleright_M \rangle$. This result is parallel to Birkhoff's Representation Theorem (Birkhoff 1937) for distributive lattices.²⁰ Next, we make some simple observations using this result. Let $\succ \in \mathcal{P}$ be a given problem with the associated set of stable matching \mathcal{S} and set of rotations \mathcal{R} . For each $\mu \in \mathcal{S}$, let R_μ be the associated closed subset of the rotation poset $\langle \mathcal{R}, \rightarrow \rangle$.

Lemma 1. *For each $\mu \in \mathcal{S}$, $\sum_{mw \in \mu} (Rank_m^A(w) + Rank_w^A(m))$ is the same.*

Proof. Let $\mu, \mu' \in \mathcal{S}$ such that $\mu' = \mu \circ \rho$ for some $\rho \in \mathcal{R}$. We first show that for each $(m, w) \in \rho$, we have

$$Rank_m^A(\mu'(m)) = Rank_m^A(w) + 1 \text{ and } Rank_w^A(\mu'(w)) = Rank_w^A(m) - 1 \quad (2)$$

To see this, note that since $\mu' = \mu \circ \rho$, for each $(m, w) \in \rho$, we have $w \succ_m \mu'(m)$

¹⁹Given two posets (S, \leq_S) and (T, \leq_T) , an *order isomorphism* from (S, \leq_S) to (T, \leq_T) is a bijective function f from S to T that is an *order embedding*, i.e. for each $x, y \in S$, $x \leq_S y$ if and only if $f(x) \leq_T f(y)$.

²⁰Asserting that for a distributive lattice L , the closed subsets of the partially ordered set induced by its *join-irreducible elements* form a distributive lattice that is isomorphic to L .

and $\mu'(w) \succ_w m$. Since there is also no other stable matching μ'' such that $\mu \triangleright_M \mu'' \triangleright_M \mu'$, it directly follows that (2) holds. It directly follows that $\sum_{mw \in \mu'} \text{Rank}_m^A(w) = \sum_{mw \in \mu} \text{Rank}_m^A(w) + |\rho|$ and $\sum_{mw \in \mu'} \text{Rank}_w^A(m) = \sum_{mw \in \mu} \text{Rank}_w^A(m) - |\rho|$. Since each $\mu \in \mathcal{S}$ can be obtained from μ^M by sequentially eliminating the rotations in $R_\mu \setminus R_{\mu^M}$, we reach the conclusion. \square

Lemma 2. *A function $F : \mathcal{S} \rightarrow \mathbb{R}$ is modular if and only if there exists an additive set function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that for each $\mu \in \mathcal{S}$, $F(\mu) = G(R_\mu)$.*

Proof. Let $R \in Cl(\mathcal{R})$. Then, it follows from Irving & Leather (1986) that there exists $\mu \in \mathcal{S}$ such that $R = R_\mu$. We define $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that for each $\mu \in \mathcal{S}$, $G(R_\mu) = F(\mu)$. Now, let $\mu, \mu' \in \mathcal{S}$. Since $\langle Cl(\mathcal{R}), \subset \rangle$ is order isomorphic to $\langle \mathcal{S}, \triangleright_M \rangle$, we have $R_{\mu \vee \mu'} = R_\mu \cap R_{\mu'}$ and $R_{\mu \wedge \mu'} = R_\mu \cup R_{\mu'}$. Then, it directly follows that F is modular if and only if $G(R_\mu \cup R_{\mu'}) = G(R_\mu) + G(R_{\mu'}) - G(R_\mu \cap R_{\mu'})$, that is G is additive. \square

Lemma 3. *Let $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ be an additive set function. Then, there exists $g : \mathcal{R} \rightarrow \mathbb{R}$ such that for each $R \in Cl(\mathcal{R})$, $G(R) = G(\emptyset) + \sum_{\rho \in R} g(\rho)$.*

Proof. Let $\rho \in \mathcal{R}$ and define its closure $C(\rho) = \{\rho' \in \mathcal{R} \mid \rho' \rightarrow \rho\}$. Since the precedence relation is transitive, $C(\rho) \setminus \{\rho\}$ is closed. Therefore, for each $\rho \in \mathcal{R}$, we can define $g(\rho) = G(C(\rho)) - G(C(\rho) \setminus \{\rho\})$. Then, for each $\rho \in \mathcal{R}$, we have $G(C(\rho)) = G(\emptyset) + \sum_{\rho' \in C(\rho)} g(\rho')$. Now, let $R \in Cl(\mathcal{R})$. Then, since $R = \bigcup_{\rho \in R} C(\rho)$, by additivity of G , we conclude that $G(R) = G(\emptyset) + \sum_{\rho \in R} g(\rho)$. \square

Lemma 4. *Let $F : \mathcal{S} \rightarrow \mathbb{R}$ be a modular function. Then, for each $i \in N$, there exists $F_i : A_i \rightarrow \mathbb{R}$ such that for each $\mu \in \mathcal{S}$, $F(\mu) = \sum_{i \in N} F_i(\mu(i))$.*

Proof. It follows from Lemma 2 and Lemma 3 that there exists $g : \mathcal{R} \rightarrow \mathbb{R}$ such that for each $\mu \in \mathcal{S}$, $F(\mu) = G(\emptyset) + \sum_{\rho \in R_\mu} g(\rho)$ for some $G(\emptyset) \in \mathbb{R}$. Now, for each $\rho \in$

\mathcal{R} and $i \in N_\rho$, define $g_i(\rho) = g(\rho)/|N_\rho|$. Note that, by construction, we have $g(\rho) = \sum_{i \in N_\rho} g_i(\rho)$. Next, for each $i \in N$ and $j \in A_i$, if i and j are matched at the men-optimal stable matching, then define $F_i(j) = G(\emptyset)/2n$. Otherwise, let ρ_{ij} be the unique rotation elimination of which makes i matched to j , and define

$$F_i(j) = \sum_{\{\rho | \rho \rightarrow \rho_{ij}\}} g_i(\rho). \quad (3)$$

Now, let $\mu \in \mathcal{S}$. Since $R_\mu \in Cl(\mathcal{R})$, we have $R_\mu = \bigcup_{i \in N} \{\rho | \rho \rightarrow \rho_{i\mu(i)}\}$. It follows that

$$\sum_{\rho \in R_\mu} g(\rho) = \sum_{i \in N} \sum_{\{\rho | \rho \rightarrow \rho_{i\mu(i)}\}} g_i(\rho). \quad (4)$$

By substituting (3) into (4), we obtain that $F(\mu) = \sum_{i \in N} F_i(\mu(i))$. \square

Next, we present a basic observation about sublattices of \mathcal{S} that is used in the proof of Corollary 1.

Lemma 5. *Let L and L' be two sublattices of \mathcal{S} such that $L \subset L'$ with the same \triangleright_M -best and \triangleright_M -worst stable matchings. Then, for each $\mu, \mu' \in L'$, if $\mu \in L$, then $\mu \vee \mu' \in L$ or $\mu \wedge \mu' \in L$.*

Proof. If $\mu' \in L$, then the conclusion follows, since L is a sublattice of \mathcal{S} . Suppose that $\mu' \notin L$. Then, by contradiction, suppose that neither $\mu \vee \mu' \in L$ nor $\mu \wedge \mu' \in L$. Assume without loss of generality that there is no other $\hat{\mu} \in L$ such that $\hat{\mu} \triangleright_M \mu$ with neither $\hat{\mu} \vee \mu' \in L$ nor $\hat{\mu} \wedge \mu' \in L$.

Now, since $L \subset L'$ with the same \triangleright_M -best and \triangleright_M -worst stable matchings, let $\mu^1, \mu^2 \in L$ be the \triangleright_M -worst and \triangleright_M -best matchings in L such that $\mu^1 \triangleright_M \mu' \triangleright_M \mu^2$.

Next, consider the matching $(\mu \vee \mu') \wedge \mu^1$. Note that L' is distributive, as a sublattice of \mathcal{S} which is distributive. Since $\mu^1 \triangleright_M \mu'$, it follows that $(\mu \vee \mu') \wedge \mu^1 = (\mu \wedge \mu^1) \vee \mu'$.

Case 1: Suppose that $(\mu \wedge \mu^1) \vee \mu' = \mu'$. Then, we show that $\mu \wedge \mu' = \mu \wedge \mu^1$, which contradicts that the latter is in L , but the former is not. To see this, when we substitute $(\mu \wedge \mu^1) \vee \mu'$ for μ' , by distributivity, we get $\mu \wedge \mu' = (\mu \wedge \mu^1) \vee (\mu \wedge \mu')$, which equals $\mu \wedge \mu^1$, since $\mu^1 \triangleright_M \mu'$.

Case 2: Suppose that $(\mu \wedge \mu^1) \vee \mu' \neq \mu'$, and let $\mu^3 = (\mu \wedge \mu^1) \vee \mu'$. Then, since $\mu^1 \triangleright_M \mu^3 \triangleright_M \mu'$, by our choice of μ^1 , it must be that $\mu^3 \notin L$. But, then consider $\mu \wedge \mu^1$. Note that $\mu \wedge \mu^1 \in L$, since L is a sublattice. Moreover, $(\mu \wedge \mu^1) \vee \mu' = \mu^3 \notin L$ and $(\mu \wedge \mu^1) \wedge \mu' = \mu \wedge \mu' \notin L$. But, this contradicts to our choice of μ , since $\mu \triangleright_M \mu \wedge \mu^1$ and $\mu \neq \mu \wedge \mu^1$ (recall that $\mu \in L$, but $\mu \wedge \mu^1 \notin L$). \square

6.2 An order isomorphism result

Let π be a stable matching rule and $\succ \in \mathcal{P}$ be a problem with the associated rotation poset $\langle \mathcal{R}, \rightarrow \rangle$ such that $\pi(\succ)$ is a sublattice of $\langle \mathcal{S}(\succ), \triangleright_M \rangle$. First, let $\bar{\mu}(\underline{\mu})$ be the \triangleright_M -best(worst) matching in $\pi(\succ)$. Then, define $\mathcal{R}^\delta = R_{\underline{\mu}} \setminus R_{\bar{\mu}}$ and

$$Cl^0(\mathcal{R}^\delta) = \{R_\mu \setminus R_{\bar{\mu}} \mid \mu \in \pi(\succ)\}.$$

Evidently, elements of $Cl^0(\mathcal{R}^\delta)$ are not closed according to the precedence relation \rightarrow , unless $\bar{\mu}$ is the men-optimal stable matching at \succ . Next, we recursively define two set collections $\{X^i\}_{i=1}^K$ and $\{\Lambda^i\}_{i=1}^K$. Then, we prove a related structural result, Proposition 3, which will be an important stepping stone in proving Theorem 1.

For $k = 1$: Consider $\min(Cl^0(\mathcal{R}^\delta), \subset)$ that consists of $R \in Cl^0(\mathcal{R}^\delta)$ such that there is no

$R' \in Cl^0(\mathcal{R}^\delta) \setminus \emptyset$ with $R' \subsetneq R$. Let $X^1 = Cl^0(\mathcal{R}^\delta)$ and $\Lambda^1 = \min(X^1, \subset)$.

For $k \geq 2$: Define $X^2 = \{R \setminus \bigcup_{R' \in \Lambda^1} R' \mid R \in X^1\}$ and $\Lambda^2 = \min(X^2, \subset)$.²¹ Similarly, for each $k \geq 1$, define $X^{k+1} = \{R \setminus \bigcup_{R' \in \Lambda^k} R' \mid R \in X^k\}$ and $\Lambda^{k+1} = \min(X^{k+1}, \subset)$.

Let $K \geq 1$ be the smallest number such that $X^{K+1} = \emptyset$. Then, $\{X^k\}_{k=1}^K$ and $\{\Lambda^k\}_{k=1}^K$ are the ordered collection of the disjoint nonempty sets that are constructed. Define $\Lambda = \bigcup_{k=1}^K \Lambda^k$. We call each member of Λ a **hyper-rotation** and generically denote by λ . Figure 6 presents a demonstration of how Λ is formed.

Lemma 6. *For each $k \in \{1, \dots, K\}$, $\langle X^k, \subset \rangle$ is a lattice.*

Proof. By induction, first, consider the case that $k = 1$, where $X^1 = Cl^0(\mathcal{R}^\delta)$. Since $\pi(\succ)$ is a sublattice of $\langle \mathcal{S}(\succ), \triangleright_M \rangle$, $Cl^0(\mathcal{R}^\delta)$ is a sublattice of $\langle Cl(\mathcal{R}^\delta), \subset \rangle$. Therefore, $\langle X^1, \subset \rangle$ is a lattice. Next, for each $k \in \{1, \dots, K-1\}$ assume that $\langle X^k, \subset \rangle$ is a lattice, and let $Q, Q' \in X^{k+1}$. By construction of X^{k+1} , there exist $R \in X^k$ and $R' \in X^k$ such that $R = Q \cup \bigcup_{\lambda \in \Lambda^k} \lambda$ and $R' = Q' \cup \bigcup_{\lambda \in \Lambda^k} \lambda$. Since $\langle X^k, \subset \rangle$ is a lattice, $R \cap R' \in X^k$ and $R \cup R' \in X^k$. Therefore, $Q \cap Q' \in X^{k+1}$ and $Q \cup Q' \in X^{k+1}$. Thus, we conclude that $\langle X^{k+1}, \subset \rangle$ is a lattice. \square

Lemma 7. *Let $\rho \in \mathcal{R}^\delta$. Then, there exists unique $\lambda_\rho \in \Lambda$ such that $\rho \in \lambda_\rho$.*

Proof. First, note that by construction, $\{\Lambda^k\}_{k=1}^K$ is collection of disjoint sets such that for each $\rho \in \mathcal{R}^\delta$, there exists $k \in \{1, \dots, K\}$ and $\lambda \in \Lambda^k$ such that $\rho \in \lambda$. Next, let $k \in \{1, \dots, K\}$ and $\lambda, \lambda' \in \Lambda^k$ be distinct. Since, by Lemma 6, $\langle X^k, \subset \rangle$ is a lattice, $\lambda \cap \lambda' \in X^k$. Then, $\Lambda^k = \min(X^k, \subset)$ implies that $\lambda \cap \lambda' = \emptyset$. \square

²¹The elements of X^2 and Λ^2 are sets of rotations that are not closed neither according to \rightarrow nor according to \Rightarrow that is to be defined later.




Figure 6: A demonstration of our construction for Λ , where $\pi(\succ)$ are the green labelled nodes.

Lemma 8. *Let $R \in Cl^0(\mathcal{R}^\delta)$. Then, $\{\lambda_\rho\}_{\rho \in R}$ partitions R .*

Proof. For each $k \in \{1, \dots, K\}$, define $\Lambda_R^k = \{\lambda \in \Lambda^k \mid \lambda \subset R\}$. Recall that $X^1 = Cl^0(\mathcal{R}^\delta)$ and since $X^{K+1} = \emptyset$, we have $X^K = \Lambda^K$. Then, since $R \in Cl^0(\mathcal{R}^\delta)$, by construction of $\{X^k\}_{k=1}^K$ and $\{\Lambda^k\}_{k=1}^K$, we have $R = \bigcup_{k=1}^K \bigcup_{\lambda \in \Lambda_R^k} \lambda$. Since for each $\rho \in R$, by Lemma 7, λ_ρ is the unique member of Λ with $\rho \in \lambda_\rho$, it follows that $R = \bigcup_{\rho \in R} \lambda_\rho$ and for each $\rho, \rho' \in R$, either $\lambda_\rho = \lambda_{\rho'}$ or $\lambda_\rho \cap \lambda_{\rho'} = \emptyset$. \square

Now, we are ready to prove Proposition 3, which we will use to prove Theorem 1.

To introduce this result, first, recall that $\Lambda = \bigcup_{k=1}^K \Lambda^k$. Then, for each distinct $\lambda, \lambda' \in \Lambda$, λ **precedes** λ' , denoted by $\lambda \Rightarrow \lambda'$, if for each $R \in Cl^0(\mathcal{R}^\delta)$, $\lambda' \subset R$ implies that $\lambda \subset R$. Put differently, $\lambda \Rightarrow \lambda'$ if

$$\lambda \subset \bigcap_{\{R \in Cl^0(\mathcal{R}^\delta) | \lambda' \subset R\}} R.$$

Note that $\langle \Lambda, \Rightarrow \rangle$ is a finite poset. Let $Cl(\Lambda)$ be the closed subsets of Λ with respect to the precedence relation \Rightarrow . In Proposition 3, we show that $\langle Cl^0(\mathcal{R}^\delta), \subset \rangle$ is *order isomorphic* to $\langle Cl(\Lambda), \subset \rangle$. To see this, define $\Lambda : Cl^0(\mathcal{R}^\delta) \rightarrow Cl(\Lambda)$ such that for each $R \in Cl^0(\mathcal{R}^\delta)$, $\Lambda(R) = \{\lambda_\rho\}_{\rho \in R}$.

Proposition 3. *The Λ mapping induces an order isomorphism between $\langle Cl^0(\mathcal{R}^\delta), \subset \rangle$ and $\langle Cl(\Lambda), \subset \rangle$.*

Proof. Let $R \in Cl^0(\mathcal{R}^\delta)$. First, we verify that $\Lambda(R) \in Cl(\Lambda)$. To see this, let $\lambda \in \Lambda(R)$ and $\lambda' \in \Lambda$ such that $\lambda' \Rightarrow \lambda$. Then, since $R \in Cl^0(\mathcal{R}^\delta)$ with $\lambda \subset R$, it follows from $\lambda' \Rightarrow \lambda$ that $\lambda' \subset R$. Therefore, $\lambda' \in \Lambda(R)$. To see that Λ is one-to-one, note that for each $R, R' \in Cl^0(\mathcal{R}^\delta)$, if $R \neq R'$ then $\Lambda(R) \neq \Lambda(R')$. To see that Λ is an order embedding, by Lemma 8, for each $R, R' \in Cl^0(\mathcal{R}^\delta)$, $\{\lambda_\rho\}_{\rho \in R}$ partitions R and $\{\lambda_{\rho'}\}_{\rho' \in R'}$ partitions R' . It follows that $R \subset R'$ if and only if $\{\lambda_\rho\}_{\rho \in R} \subset \{\lambda_{\rho'}\}_{\rho' \in R'}$.

Finally, we show that Λ is onto. To see this, let $Q \in Cl(\Lambda)$ and define $R_Q = \bigcup_{\lambda \in Q} \lambda$. We show that $R_Q \in Cl^0(\mathcal{R}^\delta)$ and $\Lambda(R_Q) = Q$. To see this, let $\lambda \in \Lambda$ and recall that, by construction, there exists $R \in Cl^0(\mathcal{R}^\delta)$ such that $\lambda \subset R$. Therefore, we can define $R_\lambda = \bigcap_{\{R \in Cl^0(\mathcal{R}^\delta) | \lambda \subset R\}} R$. Then, since $Cl^0(\mathcal{R}^\delta)$ is a lattice, we have $R_\lambda \in Cl^0(\mathcal{R}^\delta)$ and $\bigcup_{\lambda \in Q} R_\lambda \in Cl^0(\mathcal{R}^\delta)$. Next, to conclude, we show that $R_Q = \bigcup_{\lambda \in Q} R_\lambda$.

Since for each $\lambda \in Q$, $\lambda \subset R_\lambda$ and $R_Q = \bigcup_{\lambda \in Q} \lambda$, we have $R_Q \subset \bigcup_{\lambda \in Q} R_\lambda$. To see the converse, let $\lambda \in Q$ and $\rho \in R_\lambda$. We show that $\rho \in R_Q$. Since $R_\lambda \in Cl^0(\mathcal{R}^\delta)$,

by Lemma 8, $\{\lambda_{\rho'}\}_{\{\rho' \in R_\lambda\}}$ partitions R_λ . Therefore, $\lambda_\rho \subset R_\lambda$. Then, it follows from the construction of R_λ that for each $R \in Cl^0(\mathcal{R}^\delta)$, if $\lambda \subset R$ then $\lambda_\rho \subset R$, that is $\lambda_\rho \Rightarrow \lambda$. Now, since $Q \in Cl(\Lambda)$ and $\lambda \in Q$, $\lambda_\rho \Rightarrow \lambda$ implies that $\lambda_\rho \in Q$. Since $R_Q = \bigcup_{\lambda \in Q} \lambda$, we have $\lambda_\rho \subset R_Q$ indicating that $\rho \in R_Q$. Thus, we conclude that $R_Q = \bigcup_{\lambda \in Q} R_\lambda$. Then, it directly follows from the formulation of Λ that $\Lambda(R_Q) = Q$. \square

7 Appendix B

7.1 Proof of Theorem 1

Only if part: Let π be a modular stable matching rule. It directly follows from modularity that $\pi(\succ)$ is a sublattice of $\langle \mathcal{S}(\succ), \triangleright_M \rangle$. Moreover, by Lemma 2, there exists an additive set function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that $\pi(\succ) = \operatorname{argmin}_{\mu \in \mathcal{S}(\succ)} G(R_\mu)$ and by Lemma 3 there exists $g : \mathcal{R} \rightarrow \mathbb{R}$ such that for each $R \in Cl(\mathcal{R})$, $G(R) = G(\emptyset) + \sum_{\rho \in R} g(\rho)$.

To see that π satisfies *convexity*, let $\mu^*, \mu^{**} \in \pi(\succ)$ and $\mu \in \mathcal{S}(\succ)$ such that $\mu(m) \in \{\mu^*(m), \mu^{**}(m)\}$ for each $m \in M$. We show that $\mu \in \pi(\succ)$. First, let $\mu' = \mu^* \vee \mu^{**}$ and $\mu'' = \mu^* \wedge \mu^{**}$. Note that, we have $\mu(m) \in \{\mu'(m), \mu''(m)\}$ for each $m \in M$, and $\mu', \mu'' \in \pi(\succ)$ since $\pi(\succ)$ is a sublattice. Therefore, if $\mu = \mu'$ or $\mu = \mu''$ then we conclude that $\mu \in \pi(\succ)$; if not then $\mu' \triangleright_M \mu \triangleright_M \mu''$, which implies that $R_{\mu'} \subsetneq R_\mu \subsetneq R_{\mu''}$. Let $P = R_\mu \setminus R_{\mu'}$ and $P' = R_{\mu''} \setminus R_\mu$.

Next, we show that each $\rho \in P$ and each $\rho' \in P'$ are independent. To see this, recall that N_ρ denotes the set of agents in a rotation ρ . Let $\rho \in P$ and $i \in N_\rho$, then in moving from μ' to μ it must be that agent i 's mate is changed and $\mu(i) = \mu''(i)$, and in moving from μ to μ'' , i 's mate can not change, that is there is no $\rho' \in P'$ such that $i \in N_{\rho'}$. Therefore, for each $\rho \in P$ and $\rho' \in P'$, $N_\rho \cap N_{\rho'} = \emptyset$, and there is no $\rho \in P$ that immediately precedes any $\rho' \in P'$. It follows that each $\rho \in P$ and each $\rho' \in P'$ are independent.

Now, since each $\rho \in P$ and each $\rho' \in P'$ are independent, $R_{\mu'} \cup P' \in Cl(\mathcal{R})$. Let $\mu''' \in \mathcal{S}(\succ)$ be such that $R_{\mu'''} = R_{\mu'} \cup P'$. Finally, to get a contradiction that μ' minimizes G , we show that $G(R_{\mu'''}) < G(R_{\mu'})$. Since $\mu', \mu'' \in \pi(\succ)$, we have $G(R_{\mu'}) = G(R_{\mu''})$. Since

G is additive and $\{P, P'\}$ partitions $R_{\mu''} \setminus R_{\mu'}$, it follows that $\sum_{\rho \in P} g(\rho) + \sum_{\rho' \in P'} g(\rho') = 0$. Now, if $\mu \notin \pi(\succ)$, then we must have $\sum_{\rho \in P} g(\rho) > 0$, which implies that $\sum_{\rho' \in P'} g(\rho') < 0$. Then, we have $G(R_{\mu''}) < G(R_{\mu'})$. Thus, we conclude that $\mu \in \pi(\succ)$.

If part: Let π be a stable matching rule that satisfies *convexity*. Let $\succ \in \mathcal{P}$ be a problem with the associated rotation poset $\langle \mathcal{R}, \rightarrow \rangle$. To show that there exists a modular fairness measure $F : \mathcal{S}(\succ) \rightarrow \mathbb{Z}$ such that $\pi(\succ) = \operatorname{argmin}_{\mu \in \mathcal{S}(\succ)} F(\mu)$, by Lemma 2, it is sufficient to show that there exists an additive set function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that $\pi(\succ) = \operatorname{argmin}_{\mu \in \mathcal{S}(\succ)} G(R_\mu)$.

It directly follows from *convexity* of π that $\pi(\succ)$ is a sublattice of $\langle \mathcal{S}(\succ), \triangleright_M \rangle$. Therefore, Proposition 3 holds for π . In what follows, we assume that the formal objects, such as $Cl^0(\mathcal{R}^\delta)$ and Λ , defined in Section 6.2 are associated with π . Note that for each $\mu \in \mathcal{S}(\succ)$, $\mu \in \pi(\succ)$ if and only if $R_\mu \setminus R_{\bar{\mu}} \in Cl^0(\mathcal{R}^\delta)$. Therefore, to prove the result, we show that there exists an additive set function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that for each $R \in Cl(\mathcal{R})$, R minimizes G if and only if $R \setminus R_{\bar{\mu}} \in Cl^0(\mathcal{R}^\delta)$. Next, by using Proposition 3 and *convexity*, we prove the following structural result that paves the way for constructing the desired additive set function.

Lemma 9 (Partition lemma). *Let $\lambda \in \Lambda$ that contains at least two rotations and $\{P, P'\}$ be a partition of λ . Then, there exist $\rho \in P$ and $\rho' \in P'$ such that $\rho \rightarrow \rho'$ or $\rho' \rightarrow \rho$.*

Proof. By contradiction, suppose that each $\rho \in P$ and $\rho' \in P'$ are independent. Let $\lambda \in \Lambda^j$ for some $j \in \{1, \dots, K\}$. Recall that $\Lambda^j = \min(X^j, \subset)$, and consider the set $A = \bigcup_{k=1}^{j-1} \Lambda^k$, in the case that $j = 1$, assume that $A = \emptyset$. Then, we have $A \in Cl(\Lambda)$. By construction of $\langle \Lambda, \Rightarrow \rangle$, for each $\lambda' \in \Lambda$, if $\lambda' \Rightarrow \lambda$, then $\lambda' \in \Lambda^i$ for some $i < j$. Therefore, we have $A \cup \{\lambda\} \in Cl(\Lambda)$. Then, by Proposition 3, there exist $R, R' \in Cl^0(\mathcal{R}^\delta)$ such that $\Lambda(R) = A$ and $\Lambda(R') = A \cup \{\lambda\}$. Let μ and μ' be the stable matchings in $\mathcal{S}(\succ)$ associated

with R and R' , i.e. $R = R_\mu \setminus R_{\bar{\mu}}$ and $R' = R_{\mu'} \setminus R_{\bar{\mu}'}$.

First, we show that there is no $\mu'' \in \pi(\succ)$ with $\mu \triangleright_M \mu'' \triangleright_M \mu'$. Otherwise, let $R'' = R_{\mu''} \setminus R_{\bar{\mu}''}$. Since $\mu'' \in \pi(\succ)$, we have $R'' \in Cl^0(\mathcal{R}^\delta)$ and since $\mu \triangleright_M \mu'' \triangleright_M \mu'$, we have $R \subsetneq R'' \subsetneq R'$. Then, by Proposition 3, $\Lambda(R'') \in Cl(\Lambda)$ and $\Lambda(R) \subsetneq \Lambda(R'') \subsetneq \Lambda(R')$. But, since $\Lambda(R) = A$ and $\Lambda(R') = A \cup \{\lambda\}$, there can not exist such $\Lambda(R'')$, a contradiction.

Now, let μ'' be the matching obtained from μ by eliminating all the rotations in P . Since each $\rho \in P$ and $\rho' \in P'$ are independent, $N_\rho \cap N_{\rho'} = \emptyset$. Therefore, in moving from μ to μ'' it must be that if an agent i 's mate is changed, then $\mu''(i) = \mu'(i)$. It follows that for each $m \in M$, $\mu''(m) \in \{\mu(m), \mu'(m)\}$. Then, by *convexity*, we have $\mu'' \in \pi(\succ)$. But, this contradicts that there is no $\mu'' \in \pi(\succ)$ with $\mu \triangleright_M \mu'' \triangleright_M \mu'$. \square

We are now ready to construct an additive function $H : Cl(\mathcal{R}^\delta) \rightarrow \mathbb{Z}$ such that for each $R \in Cl(\mathcal{R}^\delta)$, $H(R) = 0$ if and only if $R \in Cl^0(\mathcal{R}^\delta)$. Let $\lambda \in \Lambda$ and define λ^\downarrow as the set of rotations in λ that has no successor in λ , i.e. $\lambda^\downarrow = \{q \in \lambda \mid \text{there is no } \rho' \in \lambda \text{ with } q \rightarrow \rho'\}$ (we denote a generic element of λ^\downarrow by q). Similarly define λ^\uparrow as the set of rotations in λ that has no predecessor in λ , i.e. $\lambda^\uparrow = \{\rho \in \lambda \mid \text{there is no } \rho' \in \lambda \text{ with } \rho' \rightarrow \rho\}$. Since precedence is a transitive relation and λ is a finite set, $\lambda^\downarrow \neq \emptyset$ and $\lambda^\uparrow \neq \emptyset$. If λ is not singleton, then $\lambda^\uparrow \cap \lambda^\downarrow = \emptyset$. Otherwise, suppose that there exists $\rho \in \lambda^\uparrow \cap \lambda^\downarrow$ and consider $\{p\}$ and $\lambda \setminus \{p\}$, which partitions λ . Then, there is no $\rho' \in \lambda$ such that $\rho \rightarrow \rho'$ (since $\rho \in \lambda^\downarrow$) or $\rho' \rightarrow \rho$ (since $\rho \in \lambda^\uparrow$). But, this contradicts to Lemma 9. Thus, we conclude that $\lambda^\uparrow \cap \lambda^\downarrow = \emptyset$ whenever λ is not singleton. Now, let λ^\Downarrow and λ^\Uparrow be any pair of set of rotations in λ that respectively contains λ^\downarrow and λ^\uparrow such that $\lambda^\Uparrow \cap \lambda^\Downarrow = \emptyset$.

Remark 2. For the current proof, we can assume that $\lambda^\Uparrow = \lambda^\uparrow$ and $\lambda^\Downarrow = \lambda^\downarrow$. We present this general construction foreseeing that it will be crucial in proving Theorem 2 and similar results.

Let $\lambda \in \Lambda$ that is not singleton and $\rho \in \lambda$. Then, define $\lambda^{\downarrow}(\rho)$ as the set of rotations in λ^{\downarrow} that are preceded by ρ , i.e. $\lambda^{\downarrow}(\rho) = \{q \in \lambda^{\downarrow} \mid \rho \rightarrow q\}$. Similarly, define $\lambda^{\uparrow}(\rho)$ as the set of rotations in λ^{\uparrow} that precede ρ , i.e. $\lambda^{\uparrow}(\rho) = \{\rho' \in \lambda^{\uparrow} \mid \rho' \rightarrow \rho\}$.

Next, recall that, by Lemma 7, for each $\rho \in \mathcal{R}^\delta$, there exists unique $\lambda_\rho \in \Lambda$ such that $\rho \in \lambda_\rho$. Therefore, we can define $h : \mathcal{R}^\delta \rightarrow \mathbb{Z}$ such that for each $\rho \in \mathcal{R}^\delta$, if $\lambda_\rho = \{\rho\}$, then $h(\rho) = 0$; if not, then let $\lambda = \lambda_\rho$ and define

$$h(\rho) = \begin{cases} -1 & \text{if } \rho \in \lambda^{\downarrow}, \\ \sum_{q \in \lambda^{\downarrow}(\rho)} \frac{1}{|\lambda^{\uparrow}(q)|} & \text{if } \rho \in \lambda^{\uparrow}, \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

Figure 7 demonstrates this construction. For each nonempty $Q \subset \mathcal{R}^\delta$, define $H(Q) = \sum_{\rho \in Q} h(\rho)$ and $H(\emptyset) = 0$. Note that by construction of h , for each $\lambda \in \Lambda$, $H(\lambda) = 0$. The construction of h together with Lemma 9 guarantees that the following assertion holds.




Figure 7: Values of the h function, where $\lambda^{\uparrow} = \lambda^{\uparrow} = \{\rho^1, \rho^2\}$ and $\lambda^{\downarrow} = \lambda^{\downarrow} = \{\rho^5, \rho^6\}$.

Lemma 10. Let $\lambda \in \Lambda$ and $Q \subset \lambda^\downarrow$ be a nonempty set of rotations such that $\lambda^\downarrow \setminus Q \neq \emptyset$. Then, we have $H(\bigcup_{q \in Q} \lambda^\uparrow(q)) > |Q|$.

Proof. To see this, first, let $\lambda^\uparrow(Q) = \bigcup_{q \in Q} \lambda^\uparrow(q)$ and note that, by construction of h , $H(\lambda^\uparrow(Q)) \geq |Q|$. In what follows, we show that there exists $\rho^* \in \lambda^\uparrow(Q) \cap \lambda^\uparrow$ such that $\lambda^\downarrow(\rho^*)$ contains some $q' \in \lambda^\downarrow \setminus Q$. Thus, we will conclude that $h(\rho^*) > \sum_{q \in Q \cap \lambda^\downarrow(\rho^*)} \frac{1}{|\lambda^\uparrow(q)|}$ and $H(\lambda^\uparrow(Q)) > |Q|$. To see this, let P be the set of rotations in λ that precede an element of Q . Note that $P \neq \lambda$, since $\lambda^\downarrow \setminus Q \neq \emptyset$. Then, consider P and $\lambda \setminus P$, which partitions λ . It follows from Lemma 9 that there exists $\rho \in P$ and $\rho' \in \lambda \setminus P$ such that $\rho \rightarrow \rho'$ or $\rho' \rightarrow \rho$. By our choice of P , the latter is not possible, so we have $\rho \rightarrow \rho'$. Since $\rho' \in \lambda \setminus P$, there exists $q' \in \lambda^\downarrow \setminus Q$ such that $\rho' \rightarrow q'$, and thus $\rho \rightarrow q'$. Since $\rho \in P$, there exists $\rho^* \in \lambda^\uparrow(Q) \cap \lambda^\uparrow$ such that $\rho^* \rightarrow \rho$. Thus, we conclude that $\rho^* \rightarrow q'$ as desired. \square

Next, we show that by restricting the domain of H to $Cl(\mathcal{R}^\delta)$, we will obtain the desired additive function. Figure 8 presents a demonstration of our construction.

Lemma 11. For each $R \in Cl(\mathcal{R}^\delta)$, $H(R) = 0$ if and only if $R \in Cl^0(\mathcal{R}^\delta)$.

Proof. (If part) As we noted before, by the construction of h , for each $\lambda \in \Lambda$, we have $H(\lambda) = 0$. Since, by Lemma 8, for each $R \in Cl^0(\mathcal{R}^\delta)$, $\{\lambda_\rho\}_{\rho \in R}$ partitions R , it follows that $H(R) = 0$.

(Only if part) Let $R \in Cl(\mathcal{R}^\delta)$ such that $R \notin Cl^0(\mathcal{R}^\delta)$. We show that $H(R) > 0$. First, recall that, by Lemma 7, for each $\rho \in \mathcal{R}^\delta$, there exists unique $\lambda \in \Lambda$ such that $\rho \in \lambda$. Therefore, $\{R \cap \lambda\}_{\lambda \in \Lambda}$ partitions R . Since H is additive, it follows that $H(R) = \sum_{\lambda \in \Lambda} H(R \cap \lambda)$.

Since $R \in Cl(\mathcal{R}^\delta)$ but $R \notin Cl^0(\mathcal{R}^\delta)$, by Proposition 3, there exists $\lambda \in \Lambda$, such that $R \cap \lambda \neq \emptyset$ and $\lambda \setminus R \neq \emptyset$. Therefore, λ is not singleton, and by the construction of h ,

for each $\rho \in \lambda$, $h(\rho) < 0$ ($h(\rho) > 0$) if and only if $\rho \in \lambda^\downarrow$ ($\rho \in \lambda^\uparrow$). Thus, we have $H(R \cap \lambda) = H(R \cap \lambda^\uparrow) + H(R \cap \lambda^\downarrow)$.

Since R is closed and for each $\rho \in \lambda$, either $\rho \in \lambda^\uparrow$ or ρ is preceded by a rotation in $\lambda^\uparrow \subset \lambda^\uparrow$, it follows from $R \cap \lambda \neq \emptyset$ that $R \cap \lambda^\uparrow \neq \emptyset$. Now, first, suppose that $R \cap \lambda^\downarrow = \emptyset$. Then, $H(R \cap \lambda^\downarrow) = 0$, and thus $H(R \cap \lambda) > 0$. Next, suppose that $R \cap \lambda^\downarrow \neq \emptyset$ and let $Q = R \cap \lambda^\downarrow$. Note that, since R is closed, we have $\bigcup_{q \in Q} \lambda^\uparrow(q) = R \cap \lambda^\uparrow$. Therefore, $H(R \cap \lambda) = H(\bigcup_{q \in Q} \lambda^\uparrow(q)) - |Q|$.

Finally, by Lemma 10, we will conclude that $H(R \cap \lambda) > 0$. To apply the lemma, we need to show that $\lambda^\downarrow \setminus Q \neq \emptyset$. But, if we would have $\lambda^\downarrow \subset Q$, then since R is closed and for each $\rho \in \lambda$, either $\rho \in \lambda^\downarrow$ or ρ precedes a rotation in λ^\downarrow , we must have $\lambda \subset R$, contradicting that $\lambda \setminus R \neq \emptyset$. \square

Next, we extend the domain of H from $Cl(\mathcal{R}^\delta)$ to $Cl(\mathcal{R})$. For this, first define $g : \mathcal{R} \rightarrow \mathbb{Z}$ such that for each $\rho \in \mathcal{R}$,

$$g(\rho) = \begin{cases} -1 & \text{if } \rho \in R_{\bar{\mu}}, \\ 1 & \text{if } \rho \notin R_{\bar{\mu}}, \\ h(\rho) & \text{otherwise, i.e. } \rho \in \mathcal{R}^\delta. \end{cases}$$

Then, define the additive function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ such that $G(\emptyset) = 0$ and for each nonempty $R \in Cl(\mathcal{R})$, $G(R) = \sum_{\rho \in R} g(\rho)$.

It follows from the construction of g that for each $R \in Cl(\mathcal{R})$, if R minimizes G , then $R_{\bar{\mu}} \subset R \subset R_{\underline{\mu}}$. Since $R \in Cl(\mathcal{R})$, this means that $R \setminus R_{\bar{\mu}} \in Cl(\mathcal{R}^\delta)$. Then, since for each $\rho \in \mathcal{R}^\delta$, $g(\rho) = h(\rho)$, it directly follows from Lemma 11 that for each $R \in Cl(\mathcal{R})$, R minimizes G if and only if $R \setminus R_{\bar{\mu}} \in Cl^0(\mathcal{R}^\delta)$. Thus, we complete the proof.




Figure 8: A demonstration of our construction of h and g .

7.2 Proof of Proposition 2

The *if part* was proven in the main text. To see the *only if part*, let $\succ \in \mathcal{P}$. Since π satisfies *convexity*, $\pi(\succ)$ is a sublattice of $\langle \mathcal{S}(\succ), \triangleright_M \rangle$. To see that π satisfies *IIR*, first, let $\bar{\mu}(\underline{\mu})$ be the \triangleright_M -best(worst) matching in $\pi(\succ)$. Then, recall that we obtain the men(women)-optimal stable matching for the problem \succ^π by running the men(women)-proposing Gale-Shapley algorithm, in which each man(woman) proposes to women(men) by following his(her) transformed preference list. Then, it directly follows that the men(women)-optimal stable matching for the problem \succ^π is $\bar{\mu}(\underline{\mu})$. Therefore, for each $\mu \in \mathcal{S}(\succ)$, if

$\mu \triangleright_M \bar{\mu}$ or $\underline{\mu} \triangleright_M \mu$, then $\mu \notin \mathcal{S}(\succ^\pi)$. Now, let $\mu \in \mathcal{S}(\succ)$ be such that $\bar{\mu} \triangleright_M \mu \triangleright_M \underline{\mu}$. Next, we show that if $\mu \in \mathcal{S}(\succ^\pi)$, then $\mu \in \pi(\succ)$. By contradiction, suppose that $\mu \in \mathcal{S}(\succ^\pi) \setminus \pi(\succ)$. Assume without loss of generality that there is no other $\hat{\mu} \in \mathcal{S}(\succ^\pi) \setminus \pi(\succ)$ such that $\hat{\mu} \triangleright_M \mu$.

We show that there exists $m \in M$ such that $\mu(m) \notin \pi_m(\succ)$. To see this, let $\mu', \mu'' \in \pi(\succ)$ be the \triangleright_M -worst and \triangleright_M -best matchings such that $\mu' \triangleright_M \mu \triangleright_M \mu''$. Since π satisfies *convexity* and $\mu \notin \pi(\succ)$, there exists $m \in M$ such that $\mu(m) \notin \{\mu'(m), \mu''(m)\}$. Next, we show that $\mu(m) \notin \pi_m(\succ)$. By contradiction, suppose that there exists $\tilde{\mu} \in \pi(\succ)$ with $\tilde{\mu}(m) = \mu(m)$. We show that this contradicts that there is no other $\hat{\mu} \in \mathcal{S}(\succ^\pi) \setminus \pi(\succ)$ such that $\hat{\mu} \triangleright_M \mu$. To see this, let $\mu^* = (\mu' \wedge \tilde{\mu}) \vee \mu''$. Since $\pi(\succ)$ is a lattice, $\mu^* \in \pi(\succ)$ and therefore $\mu^* \neq \mu$. Moreover, $\mu^* \neq \mu'$ and $\mu^* \neq \mu''$, since $\mu^*(m) = \mu(m)$.

Now, consider the matching $\mu \vee \mu^*$. First, we show that $\mu \vee \mu^* \neq \mu$. If not, then $\mu \triangleright_M \mu^*$. Since $\mu^* \in \pi(\succ)$ and $\mu^* \triangleright_M \mu''$, this contradicts that μ'' is the \triangleright_M -best matching in $\pi(\succ)$ such that $\mu \triangleright_M \mu''$. Next, we show that $\mu \vee \mu^* \in \mathcal{S}(\succ^\pi) \setminus \pi(\succ)$. To see this, recall that $\mu \in \mathcal{S}(\succ^\pi)$ is given and we know that $\mu^* \in \pi(\succ)$. Then, as argued in the main text, $\mu^* \in \pi(\succ)$ implies that $\mu^* \in \mathcal{S}(\succ^\pi)$. Since $\mathcal{S}(\succ^\pi)$ is a lattice, it directly follows that $\mu \vee \mu^* \in \mathcal{S}(\succ^\pi)$. To see that $\mu \vee \mu^* \notin \pi(\succ)$, recall that $\mu' \triangleright_M \mu \vee \mu^* \triangleright_M \mu$. Moreover, $\mu' \neq \mu \vee \mu^*$, since $(\mu \vee \mu^*)(m) = \mu(m)$. Then, since μ' is the \triangleright_M -worst matching in $\pi(\succ)$ such that $\mu' \triangleright_M \mu$, we must have $\mu \vee \mu^* \notin \pi(\succ)$. Thus, we conclude that $\mu \vee \mu^* \in \mathcal{S}(\succ^\pi) \setminus \pi(\succ)$ such that $\mu \vee \mu^* \triangleright_M \mu$ and $\mu \vee \mu^* \neq \mu$. But, this contradicts our choice of μ . Therefore, we conclude that $\mu(m) \notin \pi_m(\succ)$.

Now, we are ready to show that $\mu \notin \mathcal{S}(\succ^\pi)$. To see this, let $w = \underline{\mu}(m)$. Then, by the construction of \succ^π , m is the \succ_w^π -best man for w . Since $\mu(m) \notin \pi_m(\succ)$, we also have $w \succ_m^\pi \mu(m)$. Therefore, m and w form a blocking pair at μ .

7.3 Proof of Theorem 2

If part: To see that *equity undominance* is satisfied, let $\succ \in \mathcal{P}$ and let $\mu \in \mathcal{S}(\succ)$ that minimizes $-\sum_{i \in N} F_i(\mu(i))$. Since for each $i \in N$, F_i is unimodal with respect to \succ_i with mode med_i^A , it follows that there is no $\mu' \in \mathcal{S}(\succ)$ such that for each $i \in N$, $\mu'(i)$ is closer to med_i^A than $\mu(i)$.

Only if part: Let $\succ \in \mathcal{P}$ be a problem, \mathcal{S} and \mathcal{R} be the associated sets of stable matchings and rotations.

Step 1. Let $\rho = [(m_1, w_1), (m_2, w_2), \dots, (m_k, w_k)]$ be a rotation. For each $i \in \{1, \dots, k\}$, define $\rho(m_i) = w_i$ and $s_\rho(m_i) = w_{i+1}$; $\rho(w_i) = m_i$ and $s_\rho(w_i) = m_{i-1}$, where the addition and subtraction in the subscripts is modulo k . Now, for each $i \in N$ and $\rho \in \mathcal{R}$, define

$$\phi_i(\rho) = \begin{cases} 0 & \text{if there is no pair with } i \text{ in } \rho, \\ -1 & \text{if } s_\rho(i) \text{ is closer to } med_i^A \text{ than } \rho(i), \\ 1 & \text{if } \rho(i) \text{ is closer to } med_i^A \text{ than } s_\rho(i). \end{cases} \quad (5)$$

Then, since each $\mu \in \mathcal{S}$ is obtainable from μ^M by eliminating the rotations in R_μ , we have for each $i \in N$,

$$|Rank_i^A(\mu(i)) - Rank_i^A(med_i^A)| = K + \sum_{\rho \in R_\mu} \phi_i(\rho) \quad (6)$$

where $K = |Rank_i^A(\mu^M(i)) - Rank_i^A(med_i^A)| = \lfloor |A_i|/2 \rfloor - 1$.

For each $i \in N$ and $j \in A_i$, let ρ_{ij} be the unique rotation elimination of which makes i matched to j . By construction of ϕ_i , for each $\rho \in R_\mu$, if there is no pair with i , then

$\phi_i(\rho) = 0$ and if there is a pair with i , then $\rho \rightarrow \rho_{i\mu(i)}$. Therefore, we have

$$|Rank_i^A(\mu(i)) - Rank_i^A(med_i^A)| = K + \sum_{\{\rho | \rho \rightarrow \rho_{i\mu(i)}\}} \phi_i(\rho). \quad (7)$$

Step 2. By using the generality of our results in Section 7.1, we construct an additive function $G : Cl(\mathcal{R}) \rightarrow \mathbb{R}$ for π such that $G(\emptyset) = 0$ and for each nonempty $R \in Cl(\mathcal{R})$, $G(R) = \sum_{\rho \in R} g(\rho)$. To see this, let $\rho \in \mathcal{R}$ and N_ρ^+ (N_ρ^-) be the set of agents such that $\phi_i(\rho) > 0$ ($\phi_i(\rho) < 0$). First, note that for each $\rho \in \mathcal{R}$, if $N_\rho^+ = \emptyset$ ($N_\rho^- = \emptyset$), then we must have $g(\rho) < 0$ ($g(\rho) > 0$), otherwise some agent $i \in N_\rho^-$ ($i \in N_\rho^+$) must receive a positive (negative) $g_i(\rho)$ value, contradicting that the resulting F_i is unimodal. To guarantee that this is not the case, for each $\lambda \in \Lambda$ that is not singleton choose λ^\downarrow as the union of λ^\downarrow and all $\rho \in \lambda$ such that $N_\rho^+ = \emptyset$, and λ^\uparrow as the union of λ^\uparrow and all $\rho \in \lambda$ such that $N_\rho^- = \emptyset$, i.e. $\lambda^\downarrow = \lambda^\downarrow \cup \{\rho' \in \lambda \mid N_{\rho'}^+ = \emptyset\}$ and $\lambda^\uparrow = \lambda^\uparrow \cup \{\rho \in \lambda \mid N_{\rho'}^- = \emptyset\}$. Note that, we have $\lambda^\uparrow \cap \lambda^\downarrow = \emptyset$, as we know that $\lambda^\uparrow \cap \lambda^\downarrow = \emptyset$. Now, by Lemma 11, for the associated mapping H , for each $\mu \in \mathcal{S}$ with $R_{\bar{\mu}} \subset R_\mu \subset R_{\underline{\mu}}$, we have $H(R_\mu) = 0$ if and only if $\mu \in \pi(\succ)$.

In moving to G from H , there is a minor problem of directly using the construction presented at the end of Section 7.1. To fix this, we need to modify our construction of g by using the additional structure brought by *equity undominance* of π to guarantee that each F_i can be constructed as to be unimodal. To show this, let $\mu \in \pi(\succ)$ and $\rho \in \mathcal{R}$, first, we make two simple observations:

(i) If ρ is exposed in μ , then $N_\rho^+ \neq \emptyset$; otherwise let $\mu' = \mu \circ \rho$, then for each agent $i \in N$ with $\mu(i) \neq \mu'(i)$ we have $\mu'(i)$ is closer to med_i^A than $\mu(i)$, which contradicts that π satisfies *equity undominance*.

(ii) If μ is obtained from another matching μ' by eliminating ρ , i.e. $\mu = \mu' \circlearrowleft \rho$, then $N_\rho^- \neq \emptyset$; otherwise for each agent $i \in N$ with $\mu(i) \neq \mu'(i)$ we have $\mu'(i)$ is closer to med_i^A than $\mu(i)$, which contradicts that π satisfies *equity undominance*.

Now, let $\rho \notin R_{\underline{\mu}}$ ($\rho \in R_{\bar{\mu}}$). Then, as it was defined in Section 7.1, $g(\rho) = 1$ ($g(\rho) = -1$) even if $N_\rho^+ = \emptyset$ ($N_\rho^- = \emptyset$). To fix this, first, update g such that if $N_\rho^+ = \emptyset$ ($N_\rho^- = \emptyset$), then $g(\rho) = -1$ ($g(\rho) = 1$). However, following this update, it should still be the case that if $R \in Cl(\mathcal{R})$ minimizes G , then $R_{\bar{\mu}} \subset R \subset R_{\underline{\mu}}$. To guarantee this, if $\bar{\mu} \neq \mu^M$, then let $\rho \in \mathcal{R}$ be such that $\bar{\mu} = \mu' \circlearrowleft \rho$ for some $\mu \in \mathcal{S}$. By (ii), we have $N_\rho^- \neq \emptyset$. Therefore, we can pick $g(\rho)$ small enough as to guarantee that for each $R \in Cl(\mathcal{R})$, if $R \subsetneq R_{\bar{\mu}}$, then $G(R_{\bar{\mu}}) < G(R)$. Similarly, if $\underline{\mu} \neq \mu^W$, then let $\rho \in \mathcal{R}$ such that ρ is exposed in $\underline{\mu}$. By (i), we have $N_\rho^+ \neq \emptyset$. Therefore, we can pick $g(\rho)$ big enough as to guarantee that for each $R \in Cl(\mathcal{R})$, if $R_{\underline{\mu}} \subsetneq R$, then $G(R_{\underline{\mu}}) < G(R)$.

Step 3. First, for the mapping $g : \mathcal{R} \rightarrow \mathbb{R}$ that is constructed in the previous step, we show that for each $\rho \in \mathcal{R}$, there exists $\{g_i(\rho)\}_{i \in N}$ that satisfies

$$\sum_{i \in N} g_i(\rho) = g(\rho), \text{ and} \quad (8)$$

$$\text{for each } i \in N, g_i(\rho) = \alpha \phi_i(\rho) \text{ for some } \alpha > 0. \quad (9)$$

To see this, recall that g is constructed such that for each $\rho \in \mathcal{R}$, if $N_\rho^+ = \emptyset$ ($N_\rho^- = \emptyset$), then $g(\rho) < 0$ ($g(\rho) > 0$). Now, for each $i \in N$ and $\rho \in \mathcal{R}$, define $g_i(\rho)$ such that if $g(\rho) < 0$, then

$$g_i(\rho) = \begin{cases} 0 & \text{if } \phi_i(\rho) = 0, \\ \frac{(g(\rho) - |N_\rho^+|)}{|N_\rho^-|} & \text{if } \phi_i(\rho) < 0, \\ 1 & \text{if } \phi_i(\rho) > 0. \end{cases} \quad (10)$$

If $g(\rho) \geq 0$, then

$$g_i(\rho) = \begin{cases} 0 & \text{if } \phi_i(\rho) = 0, \\ -1 & \text{if } \phi_i(\rho) < 0, \\ \frac{(g(\rho) + |N_\rho^-|)}{|N_\rho^+|} & \text{if } \phi_i(\rho) > 0. \end{cases} \quad (11)$$

Note that, by this construction, $\{g_i(\rho)\}_{i \in N}$ satisfies (8) and (9). Next, recall that for each $i \in N$ and $j \in A_i$, ρ_{ij} was the unique rotation elimination of which makes i matched to j . Then, we define

$$F_i(j) = \sum_{\{\rho | \rho \rightarrow \rho_{ij}\}} g_i(\rho). \quad (12)$$

It directly follows from (7) and (9) that for each $i \in N$, F_i is unimodal with mode med_i^A . Finally, let $\mu \in \mathcal{S}$ and define $F(\mu) = \sum_{i \in N} F_i(\mu(i))$. To see that $F(\mu) = G(R_\mu)$, first, note that since $R_\mu \in Cl(\mathcal{R})$, we have $R_\mu = \bigcup_{i \in N} \{\rho | \rho \rightarrow \rho_{i\mu(i)}\}$. By (9), for each $i \in N$ and $\rho \in \mathcal{R}$, if $\phi_i(\rho) = 0$, then $g_i(\rho) = 0$. Therefore, it follows from (8) that

$$G(R_\mu) = \sum_{\rho \in R_\mu} g(\rho) = \sum_{i \in N} \sum_{\{\rho | \rho \rightarrow \rho_{i\mu(i)}\}} g_i(\rho). \quad (13)$$

By substituting (12) into (13), we conclude that $F(\mu) = G(R_\mu)$.

8 Appendix C

8.1 On the structure of (stable) mixtures

For a given fixed problem $\succ \in \mathcal{P}$, and a pair of distinct stable matchings μ' and μ'' such that $\mu' \triangleright_M \mu''$, **(women-improving-)cycles** between μ' and μ'' , denoted by $\mathcal{C}(\mu', \mu'')$, are the set of mate changes that need to be made so that μ' can be transformed into μ'' . In contrast to rotations, there can be another stable matching $\mu \in \mathcal{S}(\succ)$ such that $\mu' \triangleright_M \mu \triangleright_M \mu''$. A (women-improving) **cycle** C between μ' and μ'' is an (ordered) cyclic sequence of distinct man-woman pairs $C = [(m_1, w_1), (m_2, w_2), \dots, (m_k, w_k)]$ such that $m_i w_i \in \mu'$ and $m_i w_{i+1} \in \mu''$ for each $i \in \{1, \dots, k\}$, where the addition in the subscripts is modulo k . To **open** a cycle C , each man m_i in C is matched to w_{i+1} while all the pairs that are not in C are kept the same. For each cycle $C \in \mathcal{C}(\mu', \mu'')$, let N_C denote the set of agents that appear in cycle C . It is easy to note that for each pair of distinct cycles $C, C' \in \mathcal{C}(\mu', \mu'')$, there is no agent who appears both in C and C' , i.e. $N_C \cap N_{C'} = \emptyset$.

Lemma 12. *Let $\succ \in \mathcal{P}$ be a problem, and let $\mu', \mu'' \in \mathcal{S}(\succ)$ be a pair of distinct stable matchings such that $\mu' \triangleright_M \mu''$. Then, a matching (not necessarily stable) μ is a mixture of μ' and μ'' if and only if there is a set of cycles $\{C^k\}_{k=1}^K \subset \mathcal{C}(\mu', \mu'')$ such that μ is obtained from μ' via opening the cycles $\{C^k\}_{k=1}^K$.*

Proof. First, suppose that μ is obtained from μ' via opening cycles $\{C^k\}_{k=1}^K \subset \mathcal{C}(\mu', \mu'')$. Since for each pair of distinct cycles $C^k, C^l \in \mathcal{C}(\mu', \mu'')$, there is no agent who appears both in C^k and C^l , it directly follows that for each $m \in M$, if $\mu(m) \neq \mu'(m)$, then $\mu(m) = \mu''(m)$. Therefore, μ is a mixture of μ' and μ'' . Conversely, suppose that μ is a mixture of μ' and μ'' , and let $m \in M$ such that $\mu(m) \neq \mu'(m)$. Then, we have $\mu(m) = \mu''(m)$. By

continuing similarly we obtain the sequence $[(m, \mu'(m)), (\mu'(\mu''(m)), \mu''(m)), \dots]$, which eventually yields a cycle $C^1 \in \mathcal{C}(\mu', \mu'')$. Repeating the same construction for a man who is not in C^1 , we obtain another cycle $C^2 \in \mathcal{C}(\mu', \mu'')$. By following this procedure we obtain the desired set of cycles $\{C^k\}_{k=1}^K$. \square

Let B be a sublattice of $\mathcal{S}(\succ)$ with \triangleright_M -best matching $\bar{\mu}$ and \triangleright_M -worst matching $\underline{\mu}$. Since $\mathcal{S}(\succ)$ is a distributive lattice, B is distributive as well. Then, B is a **Boolean lattice** if B is *complemented*, i.e. for each $\mu \in B$, there exists (a complement) $\mu' \in B$ such that $\mu \vee \mu' = \bar{\mu}$ and $\mu \wedge \mu' = \underline{\mu}$.

Lemma 13. *Let $\succ \in \mathcal{P}$ be a problem, and let $\mu^1, \mu^2 \in \mathcal{S}(\succ)$ be a pair of distinct stable matchings. Then, the set of stable matchings that are mixtures of μ^1 and μ^2 is a Boolean sublattice of $\mathcal{S}(\succ)$.*

Proof. Let B be the set of stable matchings that are mixtures of μ^1 and μ^2 . Then, B is a sublattice of $\mathcal{S}(\succ)$ with \triangleright_M -best matching $\bar{\mu} = \mu^1 \vee \mu^2$ and \triangleright_M -worst matching $\underline{\mu} = \mu^1 \wedge \mu^2$. Let μ be a stable matching that is a mixture of μ^1 and μ^2 . Then, it follows from Lemma 12 that there is a set of cycles $\{C^k\}_{k=1}^K \subset \mathcal{C}(\bar{\mu}, \underline{\mu})$ such that μ is obtained from $\bar{\mu}$ via opening the cycles $\{C^k\}_{k=1}^K$. Moreover, since $\mu \in \mathcal{S}(\succ)$, R_μ is a closed set of rotations such that $R_{\bar{\mu}} \subset R_\mu \subset R_{\underline{\mu}}$. Now, let $R = R_{\bar{\mu}} \cup (R_{\underline{\mu}} \setminus R_\mu)$. Next, we argue that R is a closed set of rotations as well. To see this, it is sufficient to show that there is no $\rho \in R_\mu \setminus R_{\bar{\mu}}$ that immediately precedes any $\rho' \in R$. Now, note that an agent i appears in one of the cycles $\{C^k\}_{k=1}^K$ if and only if i appears in a rotation $\rho \in R_\mu \setminus R_{\bar{\mu}}$. Since the cycles in $\mathcal{C}(\bar{\mu}, \underline{\mu})$ are disjoint, it follows that for each $\rho \in R_\mu \setminus R_{\bar{\mu}}$ and $\rho' \in R$, $N_\rho \cap N_{\rho'} = \emptyset$. Finally, let $\mu' \in \mathcal{S}(\succ)$ be such that $R_{\mu'} = R$. It directly follows that μ' complements μ . \square

8.2 An example

We present a problem to show that several claims made throughout the main text holds. Consider the problem with eight men and women whose preferences are represented by the table in Figure 9, where each entry is associated with a man m and woman w . If m and w are attainable for each other, then the rank of w in \succ_m (the rank of m in \succ_w) is written in the bottom (top) corner. If m and w are unattainable, then the associated cells are shaded, indicating that we can freely choose the associated rank as far as it is bigger than the number of agent's total attainable mates. Note that each agent has a unique median attainable mate in this problem, the associated median attainable ranks are boxed in the table.

M \ W	a	b	c	d	w	x	y	z
1	1 3	2 5			4 2		3 4	5 1
2	4 2	3 6			2 4	5 1		1 5
3		3 4			5 1	1 3	2 5	4 2
4	5 1				1 5	4 2	3 6	2 4
5		3 1	1 3				2 2	
6		2 2		1 3			3 1	
7		1 7	5 1	4 2			3 3	2 3
8		3 3	4 2	5 1	2 3		1 7	

Figure 9: The problem.

In Figure 10, we present the associated rotations and their poset. For each agent i that appears in a rotation, the superscript $(+)$ $((-))$ means that i gets far away (closer to) from his/her attainable median. For example, $(m^+, w^-) \in \rho$ means that m gets far away from his attainable median, whereas w gets closer to her attainable median after the elimination of rotation ρ .




Figure 10: The rotation poset.

In Figure 11, we present the stable matching lattice associated with the problem in Figure 9 such that each stable matching is represented as an array $[w_1, \dots, w_8]$, where each w_i is the woman who is matched with man i . Each edge is labelled by the associated rotation whose elimination from the matching in the upper end of the edge results in the matching in the lower end of the edge. The green (lighter) colored matchings are the equity undominated ones. For this problem, we make the following observations.

1. The set of equity undominated matchings is not a sublattice of the original problem (claimed in Footnote 17). The matching $[bayxcdzw]$ is the meet of two equity undominated matchings. However, it is equity dominated by $[yabxcdzw]$. It also follows from this observation that a stable matching that is between two equity

undominated matching, according to the men-wise better than relation, can be equity dominated.

2. The unique stable matching that minimizes the total spread from the median is $[yabxcdzw]$.
3. The stable matching rule presented in Example 7, which chooses the set of attainable sex-equal stable matchings, does not satisfy *equity undominance* (claimed in Footnote 18). The unique attainable sex-equal stable matching is $\mu^* = [yxbacdzw]$, since $\sum_{m \in M} \text{Rank}_m^A(\mu^*(m)) = \sum_{w \in W} \text{Rank}_w^A(\mu^*(w)) = 22$. However, μ^* is equity dominated by the matching $[bxyacdzw]$.
4. Every mixture of stable matchings is not necessarily stable (claimed in Footnote 12). To show this, we modify the problem so that the rank of woman d for man 5 is 2 and the rank of man 5 for woman d is 3. We claim that the stable matching lattice remains unchanged after this modification. We show this by showing that this modification has no effect on the set of rotations. To see this, first note that ρ^{62} the first rotation that contains 5, and it also turns out to be the first one that contains d . Now, note that once ρ^{62} is eliminated, 5 is matched to y , where $d \succ_5 y$, and d is matched to 7, where $7 \succ_d 5$. It follows that there can be no rotation that contains $(5, d)$. Put differently, 5 and d remain unattainable for each other after the modification, and thus the rotation poset remains unchanged. Next, consider the stable matchings $\mu' = [yabxcdzw]$ and $\mu'' = [wazxybdc]$. Then let μ be the matching $[wabxydzc]$ that is obtained as a mixture of μ' and μ'' , in the sense that for each agent i , we have $\mu(i) \in \{\mu'(i), \mu''(i)\}$. Clearly, μ is not stable, since it is not stable in the original problem. Alternatively, to directly see that μ is not stable, note that $(5, d)$ forms a blocking pair in μ .




Figure 11: The associated stable matchings lattice.